


Taraxacum sect. Taraxacum

Distribution



Map info

- revised records
- unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.1-0.5**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **CR**

Life strategy (Pierce method, C-score): **55.4 %**

Life strategy (Pierce method, S-score): **0 %**

Life strategy (Pierce method, R-score): **44.7 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **rosulate**

Leaf shape: **simple - pinnately divided**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **April-June**



Flowering phase: **3 Prunus avium-Ranunculus auricomus (end of early spring)**

Flower colour: **yellow**

Flower symmetry: **zygomorphic**

Perianth type: **calyx reduced, corolla present**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **ligulate**

Calyx fusion: **pappus**

Inflorescence type: **anthodium solitarium**

Dicliny: **synoecious**

Generative reproduction type: **apomixis**

Pollination syndrome: **insect-pollination**

Pollinator spectrum: **flies s. l., nitidulids (honeybee, bumblebees, solitary bees, other Hymenoptera, hoverflies, meat flies s. l., other Diptera, butterflies, beetles, thrips, other pollinators)**

Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**

Fruit colour: **white, brown, grey**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Epilobium (mainly anemochory and autochory)**

Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**

Root metamorphosis: **primary storage root**

Storage organ: **pleiocorm, primary storage root**

Shoot life span (cyclicality): **monocyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **present**

Position of root buds: **primary root**

Role of root buds in life-history of a plant: **regenerative**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **2**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **13**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **15**

Depth of the belowground bud bank (root buds excluded) [cm]: **5**

Number of buds per shoot at the soil surface (root buds included): **2**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **20**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**

Size of the belowground bud bank (root buds included): **37**

Depth of the belowground bud bank (root buds included) [cm]: **9**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**



Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **16, 24**

Ploidy level (x): **2, 3**

2C genome size [Mbp]: **2266.19**

1Cx monoploid genome size [Mbp]: **775.43**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **5x - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **7x - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions (generalist)**

Nutrient indicator value: **7 - occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.75**

Herb layer disturbance frequency indicator value: **-0.35**

Whole-community disturbance severity indicator value: **0.59**

Herb layer disturbance severity indicator value: **0.59**

Whole-community structure based disturbance indicator value: **0.54**

Herb layer structure-based disturbance indicator value: **0.61**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **2 - optimum**

1B Siliceous cliffs and block fields: **1 - rare occurrence**

1C Walls: **2 - optimum**



1D Mobile calcareous screes: **1 - rare occurrence**

2 Alpine and subalpine grasslands



2A Alpine grasslands on siliceous bedrock: **1 - rare occurrence**

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**


3 Aquatic vegetation



- 3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**
- 4 Wetland and riverine herbaceous vegetation
- 4A Reed-beds of eutrophic still waters: **1 - rare occurrence**
- 4B Halophilous reed and sedge beds: **1 - rare occurrence**
- 4C Eutrophic vegetation of muddy substrata: **1 - rare occurrence**
- 4D Riverine reed vegetation: **1 - rare occurrence**
- 4E Reed vegetation of brooks: **1 - rare occurrence**
- 4F Mesotrophic vegetation of muddy substrata: **1 - rare occurrence**
- 4G Tall-sedge beds: **1 - rare occurrence**
- 4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**
- 4I Vegetation of nitrophilous annual hygrophilous herbs: **1 - rare occurrence**
- 4J River gravel banks: **1 - rare occurrence**
- 4K Petasites fringes of montane brooks: **1 - rare occurrence**
- 4L Nitrophilous herbaceous fringes of lowland rivers: **1 - rare occurrence**
- 5 Vegetation of springs and mires
- 5A Hard-water springs with tufa formation: **1 - rare occurrence**
- 5B Lowland to montane soft-water springs: **1 - rare occurrence**
- 5C Alpine and subalpine soft-water springs: **1 - rare occurrence**
- 5D Calcareous fens: **1 - rare occurrence**
- 5E Acidic moss-rich fens and peatland meadows: **1 - rare occurrence**
- 5F Transitional mires: **1 - rare occurrence**
- 6 Meadows and mesic pastures
- 6A Mesic Arrhenatherum meadows: **2 - optimum**
- 6B Montane mesic meadows: **2 - optimum**
- 6C Pastures and park grasslands: **2 - optimum**
- 6D Alluvial meadows of lowland rivers: **2 - optimum**
- 6E Wet Cirsium meadows: **1 - rare occurrence**
- 6F Intermittently wet Molinia meadows: **2 - optimum**
- 6G Vegetation of wet disturbed soils: **2 - optimum**
- 7 Acidophilous grasslands
- 7A Subalpine and montane acidophilous grasslands: **1 - rare occurrence**
- 7B Submontane Nardus grasslands: **1 - rare occurrence**
- 8 Dry grasslands
- 8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**
- 8B Submediterranean dry grasslands on rock outcrops: **1 - rare occurrence**
- 8C Narrow-leaved sub-continental steppes: **1 - rare occurrence**
- 8D Broad-leaved dry grasslands: **1 - rare occurrence**
- 8E Acidophilous dry grasslands: **1 - rare occurrence**
- 8F Thermophilous forest fringe vegetation: **1 - rare occurrence**
- 9 Sand grasslands and rock-outcrop vegetation
- 9B Open vegetation of acidic sands: **1 - rare occurrence**
- 9C Festuca grasslands on acidic sands: **1 - rare occurrence**
- 9D Pannonian sand steppes: **1 - rare occurrence**
- 9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**
- 9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**
- 10 Saline vegetation
- 10I Inland saline meadows: **2 - optimum**



- 10J Saline steppes: **1 - rare occurrence**
- 11 Heathlands and scrub
- 11A Dry lowland to subalpine heathlands: **1 - rare occurrence**
- 11H Subalpine deciduous scrub: **1 - rare occurrence**
- 11I Willow carrs: **1 - rare occurrence**
- 11J Willow galleries of loamy and sandy river banks: **1 - rare occurrence**
- 11L Tall mesic and xeric shrub: **1 - rare occurrence**
- 11N Low xeric scrub: **1 - rare occurrence**
- 11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**
- 12 Forests
- 12A Alder carrs: **1 - rare occurrence**
- 12B Alluvial forests: **1 - rare occurrence**
- 12C Oak-hornbeam forests: **1 - rare occurrence**
- 12D Ravine forests: **1 - rare occurrence**
- 12E Herb-rich beech forests: **1 - rare occurrence**
- 12F Limestone beech forests: **2 - optimum**
- 12G Acidophilous beech forests: **1 - rare occurrence**
- 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**
- 12I Sub-continental thermophilous oak forests: **1 - rare occurrence**
- 12J Acidophilous thermophilous oak forests: **1 - rare occurrence**
- 12K Acidophilous oak forests: **1 - rare occurrence**
- 12L Boreo-continental pine forests: **1 - rare occurrence**
- 12O Peri-Alpidic pine forests: **1 - rare occurrence**
- 12T Robinia pseudacacia plantations: **1 - rare occurrence**
- 12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**
- 12V Spruce plantations: **1 - rare occurrence**
- 12W Pine and larch plantations: **1 - rare occurrence**
- 13 Anthropogenic vegetation
- 13A Annual vegetation of ruderal habitats: **2 - optimum**
- 13B Annual vegetation of arable land: **2 - optimum**
- 13C Annual vegetation of trampled habitats: **2 - optimum**
- 13D Perennial thermophilous ruderal vegetation: **2 - optimum**
- 13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 - optimum**
- 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**
- Affinity to the forest environment
- Affinity to the forest environment in Thermophyticum: **2.2 - taxon occurring partly in the forest, but mainly in open vegetation**
- Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 - taxon occurring partly in the forest, but mainly in open vegetation**
- Constant taxon
- Constant taxon of classes: [SB Cymbalario muralis-Parietarietea judaicae](#), [TC Festuco-Puccinellietea](#), [XA Polygono arenastri-Poëtea annuae](#), [XB Stellarietea mediae](#), [XC Artemisietea vulgaris](#)
- Constant taxon of alliances: [SAA Cystopteridion](#), [SBA Cymbalario muralis-Asplenion](#), [TCB Juncion gerardii](#), [TDA Arrhenatherion elatioris](#), [TDC Cynosurion cristati](#), [TDE Deschampsion cespitosae](#), [XAA Coronopodo-Polygonion arenastri](#), [XAB Saginion procumbentis](#), [XBA Caucalidion](#), [XBB Veronico-Euphorbion](#), [XBC Scleranthion annui](#), [XBE Oxalidion fontanae](#), [XBH Sisymbrium officinalis](#), [XBK](#)



[Eragrostion cilianensi-minoris](#), [XCB Dauco carotae-Melilotion](#), [XCE Arction lappae](#), [XDD Geo urbani-Alliarion petiolatae](#)

Constant taxon of associations: [KAB01 Salicetum elaeagno-purpureae](#), [SAA01 Cystopteridetum fragilis](#), [SAA02 Asplenietum rutae-murario-trichomanis](#), [SBA01 Cymbalarietum muralis](#), [TCB01 Scorzonero parviflorae-Juncetum gerardii](#), [TCB02 Loto tenuis-Potentilletum anserinae](#), [TDA01 Pastinaco sativae-Arrhenatheretum elatioris](#), [TDA02 Ranunculo bulbosi-Arrhenatheretum elatioris](#), [TDA03 Poo-Trisetetum flavescens](#), [TDA04 Potentillo albae-Festucetum rubrae](#), [TDC01 Lolio perennis-Cynosuretum cristati](#), [TDC03 Lolietum perennis](#), [TDC04 Prunello vulgaris-Ranunculetum repentis](#), [TDE01 Poo trivialis-Alopecuretum pratensis](#), [TDE02 Holcetum lanati](#), [TDE04 Cnidio dubii-Deschampsietum cespitosae](#), [THF02 Brachypodio pinnati-Molinietum arundinaceae](#), [XAA01 Polygonetum arenastri](#), [XAA04 Eragrostio minoris-Polygonetum arenastri](#), [XAB01 Sagino procumbentis-Bryetum argentei](#), [XAB03 Rumici acetosellae-Spergularietum rubrae](#), [XAB04 Poëtum annuae](#), [XAB05 Lolio perennis-Matricarietum discoideae](#), [XBA03 Euphorbio exiguae-Melandrietum noctiflori](#), [XBA04 Stachyo annuae-Setarietum pumilae](#), [XBA05 Veronicetum hederifoliotriphylli](#), [XBB01 Mercurialietum annuae](#), [XBB02 Veronica-Lamietum hybridi](#), [XBC01 Aphano arvensis-Matricarietum chamomillae](#), [XBC03 Erophilo verna-Arabidopsietum thalianae](#), [XBE01 Echinochloo cruris-galli-Chenopodietum polyspermi](#), [XBG02 Chenopodietum urbici](#), [XBG10 Chamaeplietum officinalis](#), [XBG11 Conyzo canadensis-Lactucetum serriolae](#), [XBG12 Ivaetum xanthiifoliae](#), [XBH01 Hordeetum murini](#), [XBH02 Hordeo murini-Brometum sterilis](#), [XBI03 Polygono arenastri-Chenopodietum muralis](#), [XBI04 Malvo neglectae-Chenopodietum vulvariae](#), [XBK01 Digitario sanguinalis-Eragrostietum minoris](#), [XBK02 Portulacetum oleraceae](#), [XBK03 Eragrostio poaeoidis-Panicetum capillaris](#), [XCB01 Melilotetum albo-officinalis](#), [XCB02 Berteroetum incanae](#), [XCB03 Dauco carotae-Crepidetum rhoeadifoliae](#), [XCB04 Dauco carotae-Picridetum hieracioidis](#), [XCB05 Poo compressae-Tussilaginetum farfarae](#), [XCB06 Poëtum humili-compressae](#), [XCB07 Tanaceto vulgaris-Artemisietum vulgaris](#), [XCE01 Urtico urentis-Chenopodietum boni-henrici](#), [XCE02 Arctietum lappae](#), [XDC02 Epilobio montani-Geranium robertiani](#), [XDD01 Alliaro petiolatae-Chaerophylletum temuli](#), [XDD03 Anthriscetum trichospermae](#), [XDE01 Elytrigio repentis-Aegopodietum podagrariae](#), [XDE02 Symphyto officinalis-Anthriscetum sylvestris](#), [XDE03 Chaerophylletum aromatici](#), [XDE05 Chaerophylletum bulbosi](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **3.6**

Ecological specialization index for non-forest vegetation: **3.8**

Ecological specialization index for forest vegetation: **4**

Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional, meridional**



Floristic region: **Europe**

Distribution range extension along the continentality gradient: **7**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **669**

taxon.data.freq_in_quad: **2460**



Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **15.8 %**

Occurrence frequency in vegetation plots with a cover above 5%: **6.5 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0.8 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **2.9 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic


Number of narrow habitats in which the taxon occurs: **77**


Number of narrow habitats in which the taxon has its optimum: **16**


Number of broad habitats in which the taxon occurs: **13**


Number of broad habitats in which the taxon has its optimum: **5**











































© Vladimír Neješleba






© Vladimír Neješleba




© Vladimír Neješleba




© Vladimír Neješleba











© Vladimír Nejšchleba




© Vladimír Nejšchleba




© Vladimír Nejšchleba





© Vladimír Nejšchleba










































© Vladimír Nejšpíleba



© Vladimír Nejšpíleba





© Vladimír Nejšpíleba





© Vladimír Nejšpíleba


























































© Vladimír Neješleba



© Vladimír Neješleba



© Vladimír Neješleba





© Vladimír Neješleba






















































© Vladimír Nejeschleba



© Vladimír Nejeschleba



© Vladimír Nejeschleba



© Vladimír Nejeschleba



