
Carex pediformis agg.



Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.15–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy (Pierce method based on leaf traits): **CSR**
Life strategy (Pierce method, C-score): **36.8 %**
Life strategy (Pierce method, S-score): **30 %**
Life strategy (Pierce method, R-score): **33.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flowering period [month]: **April–May**
Flower colour: **brown**
Perianth type: **flower achlamydeous**
Inflorescence type: **spica e spiculis composita**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal



Fruit type: **dry fruit – nut enclosed in an utricle**
Fruit colour: **brown**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably myrmecochorous, probably myrmecochorous nv**

Trophic mode



Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **70**
Ploidy level (x): **4**
2C genome size [Mbp]: **1765.68**
1Cx monoploid genome size [Mbp]: **441.42**

Taxon origin



Origin in the Czech Republic: **native**

Ecological indicator values



Ellenberg-type indicator values

Light indicator value: **5 – semi-shade plant, only exceptionally occurring in full light, but usually at more than 10% of the diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **2 – optimum**
1B Siliceous cliffs and block fields: **2 – optimum**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12D Ravine forests: **1 – rare occurrence**
12L Boreo-continental pine forests: **2 – optimum**

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.8**
Ecological specialization index for forest vegetation: **4.5**

Colonization ability

Index of colonization success (ICS): **1**



Distribution and frequency

Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 52
taxon.data.freq_in_quad: 81



Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **50 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **9.3 %**
Maximum percentage cover in vegetation plots: **18 %**



Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **8**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **2**





© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz









© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz


© Dana Michalcová



