
Alchemilla effusa

Distribution



Habitus and growth type

Height [m]: **0.1–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **simple – palmately divided**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green, evergreen**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–August**
Flower colour: **yellow-green**
Perianth type: **calyx present, corolla absent**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus e cincinnis compositus**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Dicliny: **synoecious**
Generative reproduction type: **obligate apomixis**
Pollination syndrome: **insect-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Reproduction type: **by seed/spores and vegetatively, only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **rhizome, bulb**
Storage organ: **rhizome, bulb**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **2 – optimum**

4 Wetland and riverine herbaceous vegetation

4E Reed vegetation of brooks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **1 – rare occurrence**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 – rare occurrence**
5C Alpine and subalpine soft-water springs: **1 – rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **2 – optimum**
6C Pastures and park grasslands: **1 – rare occurrence**
6E Wet Cirsium meadows: **2 – optimum**

11 Heathlands and scrub

11H Subalpine deciduous scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

does not spontaneously occur in Czech forests

Distribution and frequency

Floristic zone: **southern temperate, submeridional, meridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **montane belt, subalpine belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 8
taxon.data.freq_in_quad: 11

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **9**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **2**

Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **NT – near threatened**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz