Artemisia campestris

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.3–0.8**
Growth form: **polycarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CS – competitor/stress-tolerator**
Life strategy (Pierce method based on leaf traits): **SR/CSR**
Life strategy (Pierce method, C-score): **14.7 %**
Life strategy (Pierce method, S-score): **42.5 %**
Life strategy (Pierce method, R-score): **42.9 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – pinnately divided**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **July–October**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **green-white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx absent, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **tubular, filiform**
Inflorescence type: **panicula ex anthodiis composita**
Dicliny: **gynomonoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably non-myrmecochorous**

Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**
Storage organ: **pleiocorm**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **present**
Position of root buds: **lateral roots**
Role of root buds in life-history of a plant: **additive**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **10**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **8**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **8**
Size of the belowground bud bank (root buds excluded): **25**
Depth of the belowground bud bank (root buds exluded) [cm]: **6**
Number of buds per shoot at the soil surface (root buds included): **10**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **18**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **18**
Size of the belowground bud bank (root buds included): **45**
Depth of the belowground bud bank (root buds included) [cm]: **8**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Karyology

Chromosome number (2n): **36**
Ploidy level (x): **4**
2C genome size [Mbp]: **9471.06**
1Cx monoploid genome size [Mbp]: **2367.76**
Genomic GC content: **40.8 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**
Moisture indicator value: **2 – transition between values 1 and 3**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.47**
Herb layer disturbance frequency indicator value: **-0.27**
Whole-community disturbance severity indicator value: **0.23**
Herb layer disturbance severity indicator value: **0.27**
Whole-community structure based disturbance indicator value: **0.63**
Herb layer structure-based disturbance indicator value: **0.68**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**
1B Siliceous cliffs and block fields: **1 – rare occurrence**
1D Mobile calcareous screes: **1 – rare occurrence**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 – rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8B Submediterranean dry grasslands on rock outcrops: **2 – optimum**
8C Narrow-leaved sub-continental steppes: **2 – optimum**
8D Broad-leaved dry grasslands: **1 – rare occurrence**
8E Acidophilous dry grasslands: **2 – optimum**
8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **2 – optimum**

9C Festuca grasslands on acidic sands: **2 – optimum**

9D Pannonian sand steppes: **2 – optimum**

9E Acidophilous vegetation of spring therophytes and succulents: **2 – optimum**

9F Basiphilous vegetation of spring therophytes and succulents: **2 – optimum**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**

11L Tall mesic and xeric shrub: **1 – rare occurrence**

11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**

12J Acidophilous thermophilous oak forests: **1 – rare occurrence**

12K Acidophilous oak forests: **1 – rare occurrence**

12O Peri-Alpidic pine forests: **1 – rare occurrence**

12T Robinia pseudacacia plantations: **1 – rare occurrence**

12W Pine and larch plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: **TH *Festuco-Brometea***

Diagnostic taxon of alliances: **TFC *Armerion elongatae*, THA *Alysso-Festucion pallentis*, THB *Bromo pannonici-Festucion pallentis*, THD *Festucion valesiacae***

Diagnostic taxon of associations: **TFC02 *Erysimo diffusi-Agrostietum capillaris*, THA02 *Seselio ossei-Festucetum pallentis*, THB01 *Poo badensis-Festucetum pallentis*, THD01 *Festuco valesiacae-Stipetum capillatae*, THD02 *Erysimo crepidifolii-Festucetum valesiacae*, THG02 *Avenulo pratensis-Festucetum valesiacae***

Constant taxon

Constant taxon of classes: **TG *Festucetea vaginatae***

Constant taxon of alliances: **TFC *Armerion elongatae*, TGA *Festucion vaginatae*, THA *Alysso-Festucion pallentis*, THB *Bromo pannonici-Festucion pallentis*, THD *Festucion valesiacae*, XCD *Artemisio-Kochion prostratae***

Constant taxon of associations: **TFC02 *Erysimo diffusi-Agrostietum capillaris*, TGA01 *Diantho serotini-Festucetum vaginatae*, THA02 *Seselio ossei-Festucetum pallentis*, THB01 *Poo badensis-Festucetum pallentis*, THD01 *Festuco valesiacae-Stipetum capillatae*, THD02 *Erysimo crepidifolii-Festucetum valesiacae*, THG02 *Avenulo pratensis-Festucetum valesiacae*, XCD01 *Agropyro cristati-Kochietum prostratae***

Dominant taxon

Dominant taxon of associations: **THD03 *Festuco rupicolae-Caricetum humilis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological specialization index for non-forest vegetation: **4.8**
Ecological specialization index for forest vegetation: **4.3**

Colonization ability

Index of colonization success (ICS): **7**
Index of colonization potential (ICP): **6**
Optimum successional age [years]: **32**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Western Asia**
Continentality degree: **7**
Distribution range extension along the continentality gradient: **8**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 285
taxon.data.freq_in_quad: 658

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **1.4 %**
Occurrence frequency in vegetation plots with a cover above 5%: **11.9 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.9 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **3.9 %**
Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **25**
Number of narrow habitats in which the taxon has its optimum: **9**
Number of broad habitats in which the taxon occurs: **7**
Number of broad habitats in which the taxon has its optimum: **2**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz