
Carex oederi



Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.05–0.25**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **S – stress-tolerator**
Life strategy (Pierce method based on leaf traits): **SR/CSR**
Life strategy (Pierce method, C-score): **16.4 %**
Life strategy (Pierce method, S-score): **44.4 %**
Life strategy (Pierce method, R-score): **39.3 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **evergreen**
Leaf anatomy: **mesomorphic, helomorphic**

Flower

Flowering period [month]: **June**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz









Flower colour: **green**
Perianth type: **flower achlamydeous**
Inflorescence type: **spica e spiculis composita**
Dicliny: **monoecious**
Generative reproduction type: **facultative autogamy**
Pollination syndrome: **wind-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – nut enclosed in an utricle**
Fruit colour: **green, yellow, brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably myrmecochorous**

Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome, tuft**
Type of clonal growth organ: **epigeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Karyology

Chromosome number (2n): **70**
Ploidy level (x): **2**
2C genome size [Mbp]: **695**
1Cx monoploid genome size [Mbp]: **347.5**
Genomic GC content: **35.9 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **9 – wetness indicator, focus on often soaked, poorly aerated soils**
Reaction indicator value: **7x – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions (generalist)**
Nutrient indicator value: **2 – transition between values 1 and 3**
Salinity indicator value: **3 – β-mesohaline, mostly on soils with low salt content**

Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **2 – optimum**

4 Wetland and riverine herbaceous vegetation

4H Vegetation of low annual hygrophilous herbs: **1 – rare occurrence**

5 Vegetation of springs and mires

5C Alpine and subalpine soft-water springs: **1 – rare occurrence**
5D Calcareous fens: **2 – optimum**
5E Acidic moss-rich fens and peatland meadows: **1 – rare occurrence**
5F Transitional mires: **1 – rare occurrence**
5H Wet peat soils and bog hollows: **1 – rare occurrence**

6 Meadows and mesic pastures

6G Vegetation of wet disturbed soils: **2 – optimum**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.3**
Ecological specialization index for non-forest vegetation: **4.3**

Colonization ability

Index of colonization success (ICS): **2**
Index of colonization potential (ICP): **7**
Optimum successional age [years]: **20**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional, austral or antarctic**
Floristic region: **circumpolar**
Distribution range extension along the continentality gradient: **6**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 157
taxon.data.freq_in_quad: 250

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**
Occurrence frequency in vegetation plots with a cover above 5%: **27.8 %**
Occurrence frequency in vegetation plots with a cover above 25%: **11.1 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **8.2 %**
Maximum percentage cover in vegetation plots: **42 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **9**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C2b – endangered taxon, rare and declining**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz