Carex spicata

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–1**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CS – competitor/stress-tolerator**
Life strategy (Pierce method based on leaf traits): **R/CSR**
Life strategy (Pierce method, C-score): **15.9 %**
Life strategy (Pierce method, S-score): **27.8 %**
Life strategy (Pierce method, R-score): **56.3 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flowering period [month]: **May**
Flower colour: **green**
Perianth type: **flower achlamydeous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Inflorescence type: **spica e spiculis composita**
Dicliny: **monoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **wind-pollination**




Fruit, seed and dispersal

Fruit type: **dry fruit – nut enclosed in an utricle**
Fruit colour: **brown**
Reproduction type: **mostly by seed/spores, rarely vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably non-myrmecochorous**



Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome, tuft**
Type of clonal growth organ: **epigeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **6**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **4**



Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**






Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **57**
Ploidy level (x): **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



2C genome size [Mbp]: **758.92**
1Cx monoploid genome size [Mbp]: **379.46**
Genomic GC content: **36.3 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 – taxon occurring both in the forest and open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 – taxon occurring both in the forest and open vegetation**

Colonization ability

Index of colonization success (ICS): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Western Asia**
Distribution range extension along the continentality gradient: **6**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 596
taxon.data.freq_in_quad: 1621

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz