
# *Centaurea phrygia*

## Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



### Habitus and growth type

Height [m]: **0.4–1**
Growth form: **clonal herb**
Life form: **hemicryptophyte**

### Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**

### Flower

Flowering period [month]: **July–August**
Flower colour: **pink, pink-violet**
Flower symmetry: **actinomorphic, zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **funnel-shaped, tubular**
Calyx fusion: **pappus**
Inflorescence type: **corymbothyrsus ex anthodiis compositus**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **myrmecochorous**

## Belowground organs and clonality

Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **2**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.13**
Clonal index: **4**
Position of root buds: **lateral roots**
Role of root buds in life-history of a plant: **regenerative**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **20**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**
Size of the belowground bud bank (root buds included): **40**
Depth of the belowground bud bank (root buds included) [cm]: **8**

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

**to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **4 – transition between values 3 and 5**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

## Habitat and sociology

### Occurrence in habitats

#### 4 Wetland and riverine herbaceous vegetation

4K Petasites fringes of montane brooks: **1 – rare occurrence**

#### 6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **2 – optimum**
6B Montane mesic meadows: **2 – optimum**
6C Pastures and park grasslands: **1 – rare occurrence**

#### 7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 – rare occurrence**

#### 8 Dry grasslands

8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

#### 11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 – rare occurrence**

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

### Colonization ability

Index of colonization success (ICS): **2**
Index of colonization potential (ICP): **4**
Optimum successional age [years]: **47**

## Distribution and frequency

Continentality degree: **7**
Distribution range extension along the continentality gradient: **2**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 118
taxon.data.freq_in_quad: 193

### Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **7**
Number of narrow habitats in which the taxon has its optimum: **2**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **1**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Threats and protection

Red List 2017 (national categories): **C2b – endangered taxon, rare and declining**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **not protected by law**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*