Geranium bohemicum

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.3–0.6**
Growth form: **annual herb**
Life form: **therophyte (hemicryptophyte)**
Life strategy: **CR – competitor/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite, rosulate**
Leaf shape: **simple – palmately divided**
Stipules: **present**
Petiole: **present**
Leaf life span: **overwintering green**

Flower

Flowering period [month]: **June–August**
Flower colour: **blue-violet**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Inflorescence type: **dichasium**
Dicliny: **synoecious**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – dry schizocarp with an apical beak**
Fruit colour: **brown, black**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Root metamorphosis: **primary storage root**
Storage organ: **primary storage root**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 2
taxon.data.freq_in_quad: 3

Threats and protection

Red List 2017 (national categories): **A1 – extinct taxon**
Red List 2017 (IUCN categories): **RE – regionally extinct**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz