


Hepatica nobilis

Distribution



Map info	
●	revised records
●	unrevised records
On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.	



Habitus and growth type

Height [m]: **0.05-0.15**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **CR**

Life strategy (Pierce method, C-score): **46.5 %**

Life strategy (Pierce method, S-score): **10.5 %**

Life strategy (Pierce method, R-score): **43.1 %**



Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **rosulate**

Leaf shape: **simple - palmately divided**

Stipules: **absent**

Petiole: **present**

Leaf life span: **evergreen**

Leaf anatomy: **scleromorphic, mesomorphic**



Flower

Flowering period [month]: **March-April**

Flowering phase: 1 **Corylus avellana-Leucojum vernum (pre-spring)**



Flower colour: **white, pink, blue**

Flower symmetry: **actinomorphic**

Perianth type: **homochlamydeous**

Perianth fusion: **free**

Inflorescence type: **flores solitarii**



Dicliny: **synoecious, gynomonoecious, gynodioecious**

Generative reproduction type: **mixed mating**

Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit - head of achenes**



Fruit colour: **brown**

Reproduction type: **by seed/spores and vegetatively**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **myrmecochorous**



Belowground organs and clonality

Shoot metamorphosis: **rhizome**



Storage organ: **rhizome**

Type of clonal growth organ: **epigeogenous rhizome**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicity): **monocyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **monopodial**

Primary root: **absent**

Persistence of the clonal growth organ [year]: **4**

Number of clonal offspring: **0.5**

Lateral spreading distance by clonal growth [m]: **0.03**

Clonal index: **2**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **4**



Number of buds per shoot at a depth of 0-10 cm (root buds excluded): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **19**

Depth of the belowground bud bank (root buds excluded) [cm]: **4**



Number of buds per shoot at the soil surface (root buds included): **4**

Number of buds per shoot at a depth of 0-10 cm (root buds included): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **19**

Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **14**

Ploidy level (x): **2**

2C genome size [Mbp]: **28844.28**

1Cx monoploid genome size [Mbp]: **14422.14**

Genomic GC content: **42.3 %**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **4 - transition between values 3 and 5**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.97**

Herb layer disturbance frequency indicator value: **-0.74**

Whole-community disturbance severity indicator value: **0.23**

Herb layer disturbance severity indicator value: **0.05**

Whole-community structure based disturbance indicator value: **0.07**

Herb layer structure-based disturbance indicator value: **0.17**



Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 - rare occurrence**

8 Dry grasslands

8B Submediterranean dry grasslands on rock outcrops: **1 - rare occurrence**

8D Broad-leaved dry grasslands: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 - rare occurrence**

11N Low xeric scrub: **1 - rare occurrence**

12 Forests


12B Alluvial forests: **1 - rare occurrence**

12C Oak-hornbeam forests: **2 - optimum**

12D Ravine forests: **2 - optimum**



12E Herb-rich beech forests: **2 - optimum**
 12F Limestone beech forests: **2 - optimum**
 12G Acidophilous beech forests: **1 - rare occurrence**
 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**
 12I Sub-continental thermophilous oak forests: **2 - optimum**
 12J Acidophilous thermophilous oak forests: **1 - rare occurrence**
 12K Acidophilous oak forests: **1 - rare occurrence**
 12T Robinia pseudacacia plantations: **1 - rare occurrence**
 12V Spruce plantations: **1 - rare occurrence**
 12W Pine and larch plantations: **1 - rare occurrence**



© Jitka Kimerová

13 Anthropogenic vegetation

13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.1 - taxon occurring mainly in the closed forest**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 - taxon occurring mainly in the closed forest**

Diagnostic taxon

Diagnostic taxon of classes: [**LB Carpino-Fagetea**](#), [**LC Quercetea pubescentis**](#)

Diagnostic taxon of alliances: [**LBB Carpinion betuli**](#), [**LBD Sorbo-Fagion sylvaticae**](#),
[**LBF Tilio platyphyllo-Acerion**](#), [**LCA Quercion pubescenti-petraeae**](#)

Diagnostic taxon of associations: [**KBB03 Populo tremulae-Coryletum avellanae**](#),
[**KBC04 Senecioni fuchsii-Coryletum avellanae**](#), [**LBB01 Galio sylvatici-Carpinetum betuli**](#), [**LBB02 Stellario holostaeae-Carpinetum betuli**](#), [**LBD01 Cephalanthero damasonii-Fagetum sylvaticae**](#), [**LBF01 Aceri-Tilietum**](#), [**LBF04 Seslerio albicans-Tilietum cordatae**](#), [**LCA01 Lathyro collini-Quercetum pubescentis**](#), [**LCA03 Euphorbio-Quercetum**](#)

Constant taxon

Constant taxon of alliances: [**LBB Carpinion betuli**](#), [**LBD Sorbo-Fagion sylvaticae**](#),
[**LCA Quercion pubescenti-petraeae**](#)

Constant taxon of associations: [**KBB03 Populo tremulae-Coryletum avellanae**](#),
[**KBC04 Senecioni fuchsii-Coryletum avellanae**](#), [**LBB01 Galio sylvatici-Carpinetum betuli**](#), [**LBB02 Stellario holostaeae-Carpinetum betuli**](#), [**LBD01 Cephalanthero damasonii-Fagetum sylvaticae**](#), [**LBF01 Aceri-Tilietum**](#), [**LBF04 Seslerio albicans-Tilietum cordatae**](#), [**LCA01 Lathyro collini-Quercetum pubescentis**](#), [**LCA03 Euphorbio-Quercetum**](#), [**LCC03 Melico pictae-Quercetum roboris**](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.3**

Ecological specialization index for non-forest vegetation: **4**

Ecological specialization index for forest vegetation: **5.4**

Colonization ability

Index of colonization success (ICS): **5**

Index of colonization potential (ICP): **4**

Optimum successional age [years]: **4**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Eastern Asia**

Continentality degree: **5**

Distribution range extension along the continentality gradient: **3**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 469

taxon.data.freq_in_quad: 1213

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **2.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **15.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **1 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **4.1 %**

Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **20**

Number of narrow habitats in which the taxon has its optimum: **6**

Number of broad habitats in which the taxon occurs: **5**

Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**