


# *Linaria arvensis*

## Distribution



© Vladimír Nejšlechba

### Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.1-0.5**

Growth form: **annual herb**

Life form: **therophyte**

Life strategy: **R - ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, verticillate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

## Flower

Flowering period [month]: **July-September**

Flower colour: **blue-violet**

Flower symmetry: **zygomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **personate**



© Vladimír Nejšlechba



© Vladimír Nejšlechba

Calyx fusion: **synsepalous**  
Inflorescence type: **racemus**  
Dicliny: **synoecious**  
Generative reproduction type: **autogamy**  
Pollination syndrome: **selfing**

### **Fruit, seed and dispersal**

Fruit type: **dry fruit - capsule**  
Fruit colour: **brown**  
Reproduction type: **only by seed/spores**  
Dispersal unit (diaspore): **seed**  
Dispersal strategy: **Allium (mainly autochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot life span (cyclicity): **monocyclic shoots prevailing**  
Primary root: **present**  
Position of root buds: **hypocotyl**  
Role of root buds in life-history of a plant: **necessary**  
Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
Size of the belowground bud bank (root buds excluded): **5**  
Depth of the belowground bud bank (root buds excluded) [cm]: **1**  
Number of buds per shoot at the soil surface (root buds included): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**  
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
Size of the belowground bud bank (root buds included): **5**  
Depth of the belowground bud bank (root buds included) [cm]: **1**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Taxon origin**

Origin in the Czech Republic: **archaeophyte**  
Invasion status: **casual**  
Geographic origin: **Mediterranean**  
Year of the last record: **1998**  
Period of introduction: **Late Middle Ages (1200–1500)**  
Introduction pathway: **unintentional - agriculture**

## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **7 - heat indicator, occurring in relatively warm lowlands**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

## Habitat and sociology

### Occurrence in habitats

#### 13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 - rare occurrence**

13B Annual vegetation of arable land: **1 - rare occurrence**

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

## Distribution and frequency

Floristic zone: **southern temperate, submeridional, meridional**

Floristic region: **Europe**

Continentality degree: **2**

Distribution range extension along the continentality gradient: **2**

Elevational belt in the Czech Republic: **lowlands, colline belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **211**

taxon.data.freq\_in\_quad: **307**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **2**

Number of narrow habitats in which the taxon has its optimum: **0**

Number of broad habitats in which the taxon occurs: **1**

Number of broad habitats in which the taxon has its optimum: **0**

## Threats and protection

Red List 2017 (national categories): **A2 - missing taxon**

Red List 2017 (IUCN categories): **RE - regionally extinct**

Legal protection: **not protected by law**