


# *Lotus maritimus*

## Distribution



Map info
● revised records
○ unrevised records
On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.25-0.5**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **R/CSR**

Life strategy (Pierce method, C-score): **24.3 %**

Life strategy (Pierce method, S-score): **29 %**

Life strategy (Pierce method, R-score): **46.7 %**



## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - ternate**

Stipules: **present**

Petiole: **present**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic, helomorphic**

## Flower

Flowering period [month]: **May-July**

Flower colour: **yellow**  
 Flower symmetry: **zygomorphic**  
 Perianth type: **calyx and corolla**  
 Perianth fusion: **free**  
 Calyx fusion: **synsepalous**  
 Inflorescence type: **flores solitarii**  
 Dicliny: **synoecious**  
 Pollination syndrome: **insect-pollination**



## Fruit, seed and dispersal

Fruit type: **dry fruit - legume**  
 Fruit colour: **brown, grey**  
 Reproduction type: **mostly by seed/spores, rarely vegetatively**  
 Dispersal unit (diaspore): **seed**  
 Dispersal strategy: **Allium (mainly autochory)**  
 Myrmecochory: **non-myrmecochorous (b)**



## Belowground organs and clonality

Shoot metamorphosis: **stolon, pleiocorm**  
 Root metamorphosis: **primary storage root**  
 Storage organ: **stolon, pleiocorm, primary storage root**  
 Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**  
 Branching type of stem-derived organs of clonal growth: **sympodial**  
 Primary root: **present**  
 Bud bank  
 Number of buds per shoot at the soil surface (root buds excluded):  
 Number of buds per shoot at a depth of 0-10 cm (root buds excluded):  
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded):  
 Size of the belowground bud bank (root buds excluded):  
 Depth of the belowground bud bank (root buds excluded) [cm]: **4**  
 Number of buds per shoot at the soil surface (root buds included):  
 Number of buds per shoot at a depth of 0-10 cm (root buds included):  
 Number of buds per shoot at a depth greater than 10 cm (root buds included):  
 Size of the belowground bud bank (root buds included):  
 Depth of the belowground bud bank (root buds included) [cm]: **4**



## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**  
 Carnivory: **non-carnivorous**  
 Symbiotic nitrogen fixation: **symbiosis with rhizobia**



## Karyology

Chromosome number (2n): **12**  
 Ploidy level (x): **2**  
 2C genome size [Mbp]: **1344.95**



1Cx monoploid genome size [Mbp]: **672.48**

Genomic GC content: **42 %**



## Taxon origin

Origin in the Czech Republic: **native**



## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **7 - heat indicator, occurring in relatively warm lowlands**

Moisture indicator value: **6x - transition between values 5 and 7 (generalist)**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **4 - transition between values 3 and 5**



Salinity indicator value: **3 - β-mesohaline, mostly on soils with low salt content**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.19**

Herb layer disturbance frequency indicator value: **-0.19**

Whole-community disturbance severity indicator value: **0.21**

Herb layer disturbance severity indicator value: **0.25**

Whole-community structure based disturbance indicator value: **0.52**

Herb layer structure-based disturbance indicator value: **0.6**



## Habitat and sociology

Occurrence in habitats

5 Vegetation of springs and mires

5D Calcareous fens: **2 - optimum**



6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 - rare occurrence**

6D Alluvial meadows of lowland rivers: **1 - rare occurrence**

6F Intermittently wet Molinia meadows: **1 - rare occurrence**

6G Vegetation of wet disturbed soils: **1 - rare occurrence**

8 Dry grasslands

8D Broad-leaved dry grasslands: **2 - optimum**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **2 - optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: [\*\*TC Festuco-Puccinellietea\*\*](#)

Diagnostic taxon of alliances: [\*\*TCB Juncion gerardii\*\*](#)

Diagnostic taxon of associations: [\*\*RBA05 Junco subnodulosi-Schoenetum nigricantis\*\*](#),  
[\*\*TCB01 Scorzonero parviflorae-Juncetum gerardii\*\*](#), [\*\*TCB02 Loto tenuis-\*\*](#)  
[\*\*Potentilletum anserinae\*\*](#), [\*\*THE01 Scabioso ochroleucae-Brachypodietum pinnati\*\*](#),  
[\*\*THE04 Plantagini maritimae-Caricetum flaccae\*\*](#)

Constant taxon

Constant taxon of associations: [\*\*RBA05 Junco subnodulosi-Schoenetum nigricantis\*\*](#),  
[\*\*TCB01 Scorzonero parviflorae-Juncetum gerardii\*\*](#), [\*\*THE04 Plantagini maritimae-\*\*](#)  
[\*\*Caricetum flaccae\*\*](#)



Ecological specialization indices

Ecological specialization index for all vegetation types: **4.9**

Ecological specialization index for non-forest vegetation: **4.9**

Colonization ability

Index of colonization success (ICS): **1**

Index of colonization potential (ICP): **1**



## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe**

Continentality degree: **5**

Distribution range extension along the continentality gradient: **3**

Elevational belt in the Czech Republic: **lowlands, colline belt (submontane belt)**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 118

taxon.data.freq\_in\_quad: 239

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **29.8 %**

Occurrence frequency in vegetation plots with a cover above 25%: **5.8 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **6.9 %**

Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **8**

Number of narrow habitats in which the taxon has its optimum: **3**

Number of broad habitats in which the taxon occurs: **4**

Number of broad habitats in which the taxon has its optimum: **3**



## Threats and protection

Red List 2017 (national categories): **C3 - vulnerable taxon**

Red List 2017 (IUCN categories): **NT - near threatened**

Legal protection: **not protected by law**