Maianthemum bifolium

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.05–0.25**
Growth form: **clonal herb**
Life form: **geophyte**
Life strategy: **S – stress-tolerator**
Life strategy (Pierce method based on leaf traits): **CSR**
Life strategy (Pierce method, C-score): **28.2 %**
Life strategy (Pierce method, S-score): **28.1 %**
Life strategy (Pierce method, R-score): **43.7 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **April–May**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **white**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **fleshy fruit – berry**
Fruit colour: **red**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Cornus (mainly autochory and endozoochory)**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Shoot metamorphosis: **stolon-like rhizome**
Storage organ: **stolon-like rhizome**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **1.9**
Lateral spreading distance by clonal growth [m]: **0.13**
Clonal index: **4**



Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **4**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **19**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **4**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **19**
Depth of the belowground bud bank (root buds included) [cm]: **4**





Trophic mode


Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **36 (38, 42, 54)**
Ploidy level (x): **4**
2C genome size [Mbp]: **25009.71**
1Cx monoploid genome size [Mbp]: **6252.43**
Genomic GC content: **42 %**

Taxon origin



Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **4 – transition between values 3 and 5**
Temperature indicator value: **5x – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **4 – transition between values 3 and 5**
Nutrient indicator value: **4 – transition between values 3 and 5**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.96**
Herb layer disturbance frequency indicator value: **-0.7**
Whole-community disturbance severity indicator value: **0.25**
Herb layer disturbance severity indicator value: **0.08**
Whole-community structure based disturbance indicator value: **0.08**
Herb layer structure-based disturbance indicator value: **0.16**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 – rare occurrence**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 – rare occurrence**
5C Alpine and subalpine soft-water springs: **1 – rare occurrence**
5D Calcareous fens: **1 – rare occurrence**
5E Acidic moss-rich fens and peatland meadows: **1 – rare occurrence**
5F Transitional mires: **1 – rare occurrence**
5G Raised bogs: **1 – rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **1 – rare occurrence**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **1 – rare occurrence**
7B Submontane Nardus grasslands: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

8 Dry grasslands

8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**

11H Subalpine deciduous scrub: **1 – rare occurrence**

11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12A Alder carrs: **1 – rare occurrence**

12B Alluvial forests: **2 – optimum**

12C Oak-hornbeam forests: **2 – optimum**

12D Ravine forests: **2 – optimum**

12E Herb-rich beech forests: **2 – optimum**

12F Limestone beech forests: **2 – optimum**

12G Acidophilous beech forests: **2 – optimum**

12I Sub-continental thermophilous oak forests: **1 – rare occurrence**

12K Acidophilous oak forests: **2 – optimum**

12L Boreo-continental pine forests: **1 – rare occurrence**

12Q Peatland birch forests: **1 – rare occurrence**

12R Acidophilous spruce forests: **2 – optimum**

12S Basiphilous spruce forests: **2 – optimum**

12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**

12V Spruce plantations: **2 – optimum**

12W Pine and larch plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.1 – taxon occurring mainly in the closed forest**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 – taxon occurring mainly in the closed forest**

Diagnostic taxon

Diagnostic taxon of classes: **LB *Carpino-Fagetea***

Diagnostic taxon of alliances: **LBB *Carpinion betuli***, **LBE *Luzulo-Fagion sylvaticae***

Diagnostic taxon of associations: **LBB03 *Carici pilosae-Carpinetum betuli***, **LBE02 *Calamagrostio villosae-Fagetum sylvaticae***

Constant taxon

Constant taxon of alliances: **LBB *Carpinion betuli***

Constant taxon of associations: **LBB03 *Carici pilosae-Carpinetum betuli***, **LBC03 *Carici pilosae-Fagetum sylvaticae***, **LBC04 *Athyrio distentifolii-Fagetum sylvaticae***, **LBE02 *Calamagrostio villosae-Fagetum sylvaticae***, **LBE03 *Luzulo-Abietetum albae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.9**

Ecological specialization index for non-forest vegetation: **3.6**

Ecological specialization index for forest vegetation: **5**

Colonization ability

Index of colonization success (ICS): **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Index of colonization potential (ICP): **2**
Optimum successional age [years]: **4**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Asia**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt (subalpine belt)**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 650
taxon.data.freq_in_quad: 2171

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **3.8 %**
Occurrence frequency in vegetation plots with a cover above 5%: **5.7 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0.5 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.8 %**
Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **31**
Number of narrow habitats in which the taxon has its optimum: **10**
Number of broad habitats in which the taxon occurs: **8**
Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz