


# Melilotus officinalis

## Distribution



© Dana Michalčová

### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.3-1.7**

Growth form: **monocarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CR - competitor/ruderal**

Life strategy (Pierce method based on leaf traits): **SR/CSR**

Life strategy (Pierce method, C-score): **23.5 %**

Life strategy (Pierce method, S-score): **40.1 %**

Life strategy (Pierce method, R-score): **36.4 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - ternate**

Stipules: **present**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **scleromorphic, mesomorphic**

## Flower

Flowering period [month]: **May-September**



© Dana Michalčová



© Dana Michalčová

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **yellow**

Flower symmetry: **zygomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **synsepalous**

Inflorescence type: **racemus**

Dicliny: **synoecious**

Generative reproduction type: **facultative allogamy**

Pollination syndrome: **insect-pollination, selfing**



### Fruit, seed and dispersal

Fruit type: **dry fruit - legume**

Fruit colour: **brown, grey, black**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed, fruit, infructescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**



### Belowground organs and clonality

Root metamorphosis: **primary storage root**

Storage organ: **primary storage root**

Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**

Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **10**

Depth of the belowground bud bank (root buds excluded) [cm]: **3**


Number of buds per shoot at the soil surface (root buds included): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **10**

Depth of the belowground bud bank (root buds included) [cm]: **3**



### Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**



### Karyology

Chromosome number (2n): **16**

Ploidy level (x): **2**

2C genome size [Mbp]: **2069.18**

1Cx monoploid genome size [Mbp]: **1034.59**

Genomic GC content: **38.4 %**

## Taxon origin

Origin in the Czech Republic: **archaeophyte**

Invasion status: **naturalized**

Geographic origin: **Mediterranean, Asia**

Period of introduction: **Bronze Age (2300-750 BCE)**

Introduction pathway: **intentional - other, unintentional - anthropogenic**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.23**

Herb layer disturbance frequency indicator value: **-0.23**

Whole-community disturbance severity indicator value: **0.64**

Herb layer disturbance severity indicator value: **0.65**

Whole-community structure based disturbance indicator value: **0.58**

Herb layer structure-based disturbance indicator value: **0.66**

## Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1D Mobile calcareous screes: **1 - rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 - rare occurrence**

6G Vegetation of wet disturbed soils: **1 - rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**

8B Submediterranean dry grasslands on rock outcrops: **1 - rare occurrence**

8C Narrow-leaved sub-continental steppes: **1 - rare occurrence**

8D Broad-leaved dry grasslands: **2 - optimum**

8E Acidophilous dry grasslands: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 - rare occurrence**  
9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**  
9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**  
10 Saline vegetation  
10I Inland saline meadows: **1 - rare occurrence**  
11 Heathlands and scrub  
11L Tall mesic and xeric shrub: **1 - rare occurrence**  
11R Scrub and pioneer woodland of forests clearings: **2 - optimum**  
12 Forests  
12H Peri-Alpidic basiphilous thermophilous oak forests: **1 - rare occurrence**  
12J Acidophilous thermophilous oak forests: **1 - rare occurrence**  
12T Robinia pseudacacia plantations: **1 - rare occurrence**  
12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**  
13 Anthropogenic vegetation  
13A Annual vegetation of ruderal habitats: **2 - optimum**  
13B Annual vegetation of arable land: **1 - rare occurrence**  
13C Annual vegetation of trampled habitats: **1 - rare occurrence**  
13D Perennial thermophilous ruderal vegetation: **3 - dominant**  
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**  
Affinity to the forest environment  
Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**  
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**  
Diagnostic taxon  
Diagnostic taxon of alliances: [XCA Onopordion acanthii](#)  
Diagnostic taxon of associations: [XCA01 Carduo acanthoidis-Onopordetum acanthii](#),  
[XCB01 Melilotetum albo-officinalis](#), [XCB04 Dauco carotae-Picridetum hieracioidis](#)  
Constant taxon  
Constant taxon of associations: [XCB01 Melilotetum albo-officinalis](#)  
Dominant taxon  
Dominant taxon of associations: [XCB01 Melilotetum albo-officinalis](#)  
Ecological specialization indices  
Ecological specialization index for all vegetation types: **4.3**  
Ecological specialization index for non-forest vegetation: **4.2**  
Colonization ability  
Index of colonization success (ICS): **6**  
Index of colonization potential (ICP): **8**  
Optimum successional age [years]: **23**

## Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe, Western Asia**

Continental degree: **8**

Distribution range extension along the continentality gradient: **6**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 536

taxon.data.freq\_in\_quad: 1274

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.9 %**

Occurrence frequency in vegetation plots with a cover above 5%: **18.3 %**

Occurrence frequency in vegetation plots with a cover above 25%: **8.8 %**

Occurrence frequency in vegetation plots with a cover above 50%: **5.5 %**

Mean percentage cover in vegetation plots: **8.1 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **24**

Number of narrow habitats in which the taxon has its optimum: **4**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **3**

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**