


# Noccaea montana

## Distribution




### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.1-0.2**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CS - competitor/stress-tolerator**

Life strategy (Pierce method based on leaf traits): **R**

Life strategy (Pierce method, C-score): **12.6 %**

Life strategy (Pierce method, S-score): **0 %**

Life strategy (Pierce method, R-score): **87.4 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, rosulate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **both present and absent**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **April-May**

Flowering phase: **3 Prunus avium-Ranunculus auricomus (end of early spring)**

Flower colour: **white**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **aposepalous**

Inflorescence type: **racemus**

Dicliny: **synoecious**

Generative reproduction type: **facultative autogamy**

Pollination syndrome: **insect-pollination, selfing**



## Fruit, seed and dispersal

Fruit type: **dry fruit - silicula**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**

## Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**

Storage organ: **pleiocorm**

Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **present**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **20**

Depth of the belowground bud bank (root buds excluded) [cm]: **4**

Number of buds per shoot at the soil surface (root buds included): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **20**

Depth of the belowground bud bank (root buds included) [cm]: **4**

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **28**

Ploidy level (x): **4**

2C genome size [Mbp]: **1025.53**

1Cx monoploid genome size [Mbp]: **256.38**

Genomic GC content: **41.3 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.79**

Herb layer disturbance frequency indicator value: **-0.75**

Whole-community disturbance severity indicator value: **0.21**

Herb layer disturbance severity indicator value: **0.11**

Whole-community structure based disturbance indicator value: **0.21**

Herb layer structure-based disturbance indicator value: **0.26**

## Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8B Submediterranean dry grasslands on rock outcrops: **2 - optimum**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 - rare occurrence**

11N Low xeric scrub: **1 - rare occurrence**

12 Forests

12D Ravine forests: **1 - rare occurrence**

12F Limestone beech forests: **1 - rare occurrence**

12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**

12L Boreo-continental pine forests: **1 - rare occurrence**

12O Peri-Alpidic pine forests: **2 - optimum**

12T Robinia pseudacacia plantations: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon**

**occurring both in the forest and open vegetation**

## Diagnostic taxon

Diagnostic taxon of classes: [LE \*Erico-Pinetea\*](#)Diagnostic taxon of alliances: [LCA \*Quercion pubescenti-petraeae\*](#), [LEA \*Erico carneae-Pinion\*](#)Diagnostic taxon of associations: [LBF04 \*Seslerio albicantis-Tilietum cordatae\*](#), [LCA01 \*Lathyro collini-Quercetum pubescentis\*](#), [LEA01 \*Thlaspio montani-Pinetum sylvestris\*](#), [THC04 \*Asplenio cuneifolii-Seslerietum caeruleae\*](#)

## Constant taxon

Constant taxon of classes: [LE \*Erico-Pinetea\*](#)Constant taxon of alliances: [LEA \*Erico carneae-Pinion\*](#)Constant taxon of associations: [LEA01 \*Thlaspio montani-Pinetum sylvestris\*](#), [THC04 \*Asplenio cuneifolii-Seslerietum caeruleae\*](#)

## Ecological specialization indices

Ecological specialization index for all vegetation types: **4.7**Ecological specialization index for non-forest vegetation: **4.6**Ecological specialization index for forest vegetation: **5.2**

## Colonization ability

Index of colonization success (ICS): **2**Index of colonization potential (ICP): **5**Optimum successional age [years]: **75****Distribution and frequency**Floristic zone: **southern temperate, submeridional**Floristic region: **Europe**Continentality degree: **4**Distribution range extension along the continentality gradient: **2**Elevational belt in the Czech Republic: **colline belt**Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **18**taxon.data.freq\_in\_quad: **28**

## Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**Occurrence frequency in vegetation plots with a cover above 5%: **23.1 %**Occurrence frequency in vegetation plots with a cover above 25%: **0 %**Occurrence frequency in vegetation plots with a cover above 50%: **0 %**Mean percentage cover in vegetation plots: **5 %**Maximum percentage cover in vegetation plots: **23 %**

## Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **10**Number of narrow habitats in which the taxon has its optimum: **3**Number of broad habitats in which the taxon occurs: **3**Number of broad habitats in which the taxon has its optimum: **2****Threats and protection**Red List 2017 (national categories): **C3 - vulnerable taxon**

Red List 2017 (IUCN categories): **NT - near threatened**

Legal protection: **not protected by law**