


Petroselinum crispum

Distribution



© Dana Michalčová

Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.4-0.8**Growth form: **monocarpic perennial non-clonal herb**Life form: **hemicryptophyte**Life strategy: **CR - competitor/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**Leaf arrangement (phyllotaxis): **alternate**Leaf shape: **compound - bipinnate, compound - tripinnate**Stipules: **absent**Petiole: **both present and absent**Leaf life span: **evergreen**Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June-July**Flower colour: **green-white**Flower symmetry: **actinomorphic**Perianth type: **calyx absent, corolla present**Perianth fusion: **free**

© Vladimír Motýčka



© Vladimír Motýčka

Inflorescence type: **umbella composita**
Dicliny: **andromonoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - cremocarp**
Fruit colour: **brown, grey**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Root metamorphosis: **primary storage root**
Storage organ: **primary storage root**
Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**
Primary root: **present**
Bud bank
Number of buds per shoot at the soil surface (root buds excluded):
Number of buds per shoot at a depth of 0–10 cm (root buds excluded):
Number of buds per shoot at a depth greater than 10 cm (root buds excluded):
Size of the belowground bud bank (root buds excluded):
Depth of the belowground bud bank (root buds excluded) [cm]:
Number of buds per shoot at the soil surface (root buds included):
Number of buds per shoot at a depth of 0–10 cm (root buds included):
Number of buds per shoot at a depth greater than 10 cm (root buds included):
Size of the belowground bud bank (root buds included):
Depth of the belowground bud bank (root buds included) [cm]:

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **casual**
Geographic origin: **Mediterranean**
Period of introduction: **Late Middle Ages (1200–1500)**
Introduction pathway: **intentional - crops**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **meridional**

Floristic region: **Europe, Africa**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 32

taxon.data.freq_in_quad: 36

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**

Occurrence frequency in vegetation plots with a cover above 5%: **100 %**

Occurrence frequency in vegetation plots with a cover above 25%: **50 %**

Occurrence frequency in vegetation plots with a cover above 50%: **50 %**

Mean percentage cover in vegetation plots: **50.5 %**

Maximum percentage cover in vegetation plots: **88 %**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**