


# *Phlox paniculata*

## Distribution



© Dana Michalcová

### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Dana Holubová



© Dana Holubová



© Marek Chytrý

## Habitus and growth type

Height [m]: **0.4-1.2**

Life form: **hemicryptophyte (geophyte)**

Life strategy: **C - competitor**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, opposite**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **July-September**

Flower colour: **white, pink, red, violet**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **hypocrateriform**

Calyx fusion: **synsepalous**  
 Inflorescence type: **panicula e dichasiis composita**  
 Dicliny: **synoecious**  
 Generative reproduction type: **mixed mating**  
 Pollination syndrome: **insect-pollination**

### Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**  
 Reproduction type: **by seed/spores and vegetatively**  
 Dispersal strategy: **Zea (no dispersal)**  
 Myrmecochory: **non-myrmecochorous (b)**

### Belowground organs and clonality

Shoot metamorphosis: **rhizome**  
 Storage organ: **rhizome**

### Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**  
 Carnivory: **non-carnivorous**  
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin


Origin in the Czech Republic: **neophyte**  
 Invasion status: **casual**  
 Geographic origin: **North America**  
 Year of the first record in the wild: **1880**  
 Period of introduction: **Late Modern Period (1800-1950)**  
 Introduction pathway: **intentional - ornamental**

### Habitat and sociology

Affinity to the forest environment  
 Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**  
 Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

### Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**  
 Floristic region: **Eastern America**  
 Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 40  
 taxon.data.freq\_in\_quad: 48



## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**



© Dana Michalčová