
Pilosella bifurca



Distribution



Map info

- *revised records*
- *unrevised records*

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.1–0.45**
Growth form: **clonal herb**
Life form: **hemicryptophyte**



Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate, rosulate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**



Flower

Flowering period [month]: **May–June**
Flower colour: **yellow**
Flower symmetry: **zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **ligulate**
Calyx fusion: **pappus**
Inflorescence type: **dichasium ex anthodiis compositum**



© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Dicliny: **synoecious**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon, rhizome**
Storage organ: **rhizome**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8C Narrow-leaved sub-continental steppes: **2 – optimum**
8E Acidophilous dry grasslands: **1 – rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 – rare occurrence**
9C Festuca grasslands on acidic sands: **2 – optimum**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **2 – optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 18

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

taxon.data.freq_in_quad: 24

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **3**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz