
Pilosella visianii

Distribution



Habitus and growth type

Height [m]: **0.15–0.5**
Growth form: **clonal herb**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf shape: **simple – entire**
Stipules: **absent**

Flower

Flowering period [month]: **June**
Flower colour: **yellow**
Flower symmetry: **zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **ligulate**
Calyx fusion: **pappus**
Inflorescence type: **corymbothyrsus ex anthodiis compositus**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 4
taxon.data.freq_in_quad: 5

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz