


Ranunculus repens

Distribution



© Pavel Veselý

Map info


revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Dana Michalová



© Alena Lepší



© Pavel Veselý

Habitus and growth type

Height [m]: **0.1-0.4**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **CR**

Life strategy (Pierce method, C-score): **53 %**

Life strategy (Pierce method, S-score): **8 %**

Life strategy (Pierce method, R-score): **39.1 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - ternate**

Stipules: **absent**

Petiole: **both present and absent**

Leaf life span: **evergreen**

Leaf anatomy: **hygromorphic, helomorphic**

Flower

Flowering period [month]: **May-August**

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**

Flower colour: **yellow**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **aposepalous**

Inflorescence type: **flores solitarii**

Dicliny: **synoecious, gynodioecious**

Generative reproduction type: **allogamy self-incompatibility**

Pollination syndrome: **insect-pollination, selfing**

Pollinator spectrum: **hoverflies, flies s. l., other Diptera (honeybee, bumblebees, solitary bees, other Hymenoptera, meat flies s. l., butterflies, beetles, nitidulids, thrips, other pollinators)**

Fruit, seed and dispersal

Fruit type: **dry fruit - head of achenes**

Fruit colour: **brown**

Reproduction type: **by seed/spores and vegetatively**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon**

Storage organ: **stolon**

Type of clonal growth organ: **stolon**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **absent**

Persistence of the clonal growth organ [year]: **1.3**

Number of clonal offspring: **5.1**

Lateral spreading distance by clonal growth [m]: **0.25**

Clonal index: **6**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **12**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **8**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **20**

Depth of the belowground bud bank (root buds excluded) [cm]: **3**


Number of buds per shoot at the soil surface (root buds included): **12**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **8**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **20**

Depth of the belowground bud bank (root buds included) [cm]: **3**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **32 (16)**

Ploidy level (x): **4 (2)**

2C genome size [Mbp]: **20046.15**

1Cx monoploid genome size [Mbp]: **5011.54**

Genomic GC content: **44.4 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **5x - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

Moisture indicator value: **7 - humidity indicator, focus on well moistened, but not wet soils**

Reaction indicator value: **7x - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions (generalist)**

Nutrient indicator value: **7 - occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.94**

Herb layer disturbance frequency indicator value: **-0.4**

Whole-community disturbance severity indicator value: **0.44**

Herb layer disturbance severity indicator value: **0.45**

Whole-community structure based disturbance indicator value: **0.49**

Herb layer structure-based disturbance indicator value: **0.58**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**


3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation



- 4A Reed-beds of eutrophic still waters: **1 - rare occurrence**
- 4B Halophilous reed and sedge beds: **1 - rare occurrence**
- 4C Eutrophic vegetation of muddy substrata: **1 - rare occurrence**
- 4D Riverine reed vegetation: **2 - optimum**
- 4E Reed vegetation of brooks: **2 - optimum**
- 4F Mesotrophic vegetation of muddy substrata: **1 - rare occurrence**
- 4G Tall-sedge beds: **1 - rare occurrence**
- 4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**
- 4I Vegetation of nitrophilous annual hygrophilous herbs: **2 - optimum**
- 4J River gravel banks: **2 - optimum**
- 4K Petasites fringes of montane brooks: **2 - optimum**
- 4L Nitrophilous herbaceous fringes of lowland rivers: **1 - rare occurrence**
- 5 Vegetation of springs and mires
- 5A Hard-water springs with tufa formation: **1 - rare occurrence**
- 5B Lowland to montane soft-water springs: **2 - optimum**
- 5D Calcareous fens: **1 - rare occurrence**
- 5E Acidic moss-rich fens and peatland meadows: **1 - rare occurrence**
- 5F Transitional mires: **1 - rare occurrence**
- 6 Meadows and mesic pastures
- 6A Mesic Arrhenatherum meadows: **2 - optimum**
- 6B Montane mesic meadows: **1 - rare occurrence**
- 6C Pastures and park grasslands: **2 - optimum**
- 6D Alluvial meadows of lowland rivers: **2 - optimum**
- 6E Wet Cirsium meadows: **2 - optimum**
- 6F Intermittently wet Molinia meadows: **2 - optimum**
- 6G Vegetation of wet disturbed soils: **2 - optimum**
- 7 Acidophilous grasslands
- 7B Submontane Nardus grasslands: **1 - rare occurrence**
- 8 Dry grasslands
- 8F Thermophilous forest fringe vegetation: **1 - rare occurrence**
- 9 Sand grasslands and rock-outcrop vegetation
- 9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**
- 10 Saline vegetation
- 10I Inland saline meadows: **2 - optimum**
- 11 Heathlands and scrub
- 11H Subalpine deciduous scrub: **1 - rare occurrence**
- 11I Willow carrs: **1 - rare occurrence**
- 11J Willow galleries of loamy and sandy river banks: **2 - optimum**
- 11L Tall mesic and xeric shrub: **1 - rare occurrence**
- 11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**
- 12 Forests
- 12A Alder carrs: **2 - optimum**
- 12B Alluvial forests: **2 - optimum**
- 12C Oak-hornbeam forests: **1 - rare occurrence**
- 12D Ravine forests: **1 - rare occurrence**
- 12E Herb-rich beech forests: **1 - rare occurrence**



- 12F Limestone beech forests: **1 - rare occurrence**
 12G Acidophilous beech forests: **1 - rare occurrence**
 12I Sub-continental thermophilous oak forests: **1 - rare occurrence**
 12K Acidophilous oak forests: **1 - rare occurrence**
 12R Acidophilous spruce forests: **1 - rare occurrence**
 12T Robinia pseudacacia plantations: **1 - rare occurrence**
 12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**
 12V Spruce plantations: **1 - rare occurrence**
 12W Pine and larch plantations: **1 - rare occurrence**
 13 Anthropogenic vegetation
 13A Annual vegetation of ruderal habitats: **1 - rare occurrence**
 13B Annual vegetation of arable land: **2 - optimum**
 13C Annual vegetation of trampled habitats: **1 - rare occurrence**
 13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**
 13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 - optimum**
 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Constant taxon

Constant taxon of classes: [TC *Festuco-Puccinellietea*](#)

Constant taxon of alliances: [RAA *Caricion remotae*](#), [TCB *Juncion gerardii*](#), [TDE *Deschampsion cespitosae*](#), [XDB *Petasition hybridi*](#), [XDF *Rumicion alpini*](#)

Constant taxon of associations: [KAB02 *Salicetum purpureae*](#), [LAA03 *Carici acutiformis-Alnetum glutinosae*](#), [LBA01 *Alnetum incanae*](#), [LBA02 *Piceo abietis-Alnetum glutinosae*](#), [LBA03 *Carici remotae-Fraxinetum excelsioris*](#), [MAC03 *Pulicario vulgaris-Menthetum pulegii*](#), [MCD03 *Tussilagini farfarae-Calamagrostietum pseudophragmitae*](#), [MCE02 *Glycerietum notatae*](#), [MCG04 *Comaro palustris-Caricetum cespitosae*](#), [MCH05 *Caricetum distichae*](#), [MCH07 *Caricetum vulpinae*](#), [RAA01 *Caricetum remotae*](#), [RAA02 *Cardamino-Chrysosplenietum alternifolii*](#), [RBA03 *Valeriano simplicifoliae-Caricetum flavae*](#), [TCB01 *Scorzonero parviflorae-Juncetum gerardii*](#), [TCB02 *Loto tenuis-Potentilletum anserinae*](#), [TCB03 *Agrostio stoloniferae-Juncetum ranarii*](#), [TDC01 *Lolio perennis-Cynosuretum cristati*](#), [TDC05 *Alchemillo hybridae-Poëtum supinae*](#), [TDE01 *Poo trivialis-Alopecuretum pratensis*](#), [TDE02 *Holcetum lanati*](#), [TDE03 *Lathyro palustris-Gratioletum officinalis*](#), [TDE04 *Cnidio dubii-Deschampsietum cespitosae*](#), [TDF01 *Angelico sylvestris-Cirsietum oleracei*](#), [TDF02 *Cirsietum rivularis*](#), [TDF04 *Crepido paludosae-Juncetum acutiflori*](#), [TDF07 *Scirpo sylvatici-Cirsietum cani*](#), [TDF08 *Scirpetum sylvatici*](#), [TDF09 *Caricetum cespitosae*](#), [TDF10 *Scirpo sylvatici-Caricetum brizoidis*](#), [TDF11 *Junco inflexi-Menthetum longifoliae*](#), [XBC02 *Spergulo arvensis-Scleranthetum annui*](#), [XDB02 *Petasitetum hybrido-kablikiani*](#), [XDC01 *Stachyo sylvaticae-Impatientetum nolitangere*](#), [XDC04 *Carici pendulae-Eupatorietum cannabini*](#), [XDE03 *Chaerophylletum aromatici*](#), [XDE06 *Anthriscio nitidae-Aegopodietum podagrariae*](#), [XDF01 *Rumicetum alpini*](#)

Dominant taxon

Dominant taxon of associations: [MCC12 *Tripleurospermo inodori-Bolboschoenetum planiculmis*](#), [MCH07 *Caricetum vulpinae*](#), [TDE01 *Poo trivialis-Alopecuretum*](#)

[pratensis](#), [TDE02 Holcetum lanati](#), [TDF07 Scirpo sylvatici-Cirsietum cani](#), [TDF09 Caricetum cespitosae](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **3.8**

Ecological specialization index for non-forest vegetation: **4**

Ecological specialization index for forest vegetation: **5**

Colonization ability

Index of colonization success (ICS): **8**

Index of colonization potential (ICP): **4**

Optimum successional age [years]: **12**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe, Asia**

Distribution range extension along the continentality gradient: **7**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt (subalpine belt)**

Expansive taxon in the region: **Bohemian Thermophyticum, Bohemian Moravian Mesophyticum, Bohemian Moravian Oreophyticum, Pannonian Thermophyticum, Carpathian Mesophyticum, Carpathian Oreophyticum**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **672**

taxon.data.freq_in_quad: **2489**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **11.7 %**

Occurrence frequency in vegetation plots with a cover above 5%: **13.3 %**

Occurrence frequency in vegetation plots with a cover above 25%: **2.9 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0.5 %**

Mean percentage cover in vegetation plots: **4.5 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **55**

Number of narrow habitats in which the taxon has its optimum: **18**

Number of broad habitats in which the taxon occurs: **12**

Number of broad habitats in which the taxon has its optimum: **7**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**