


# *Rhamnus cathartica*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **1.5-4**

Growth form: **shrub**

Life form: **nanophanerophyte**

Life strategy: **C - competitor**

Life strategy (Pierce method based on leaf traits): **CSR**

Life strategy (Pierce method, C-score): **27.4 %**

Life strategy (Pierce method, S-score): **42 %**

Life strategy (Pierce method, R-score): **30.6 %**



## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **opposite**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **present**

Leaf life span: **summer green**

Leaf deciduousness in woody plants: **winter deciduous**

Leaf anatomy: **mesomorphic**

Functional leaf type in woody plants: **broad deciduous or semi-deciduous**



## Flower

Flowering period [month]: **May-June**

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **yellow-green**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **rotate**

Calyx fusion: **synsepalous**

Inflorescence type: **fasciculus**

Dicliny: **dioecious**

Generative reproduction type: **allogamy**

Pollination syndrome: **insect-pollination**



## Fruit, seed and dispersal

Fruit type: **fleshy fruit - drupe**

Fruit colour: **black**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**

Dispersal strategy: **Cornus (mainly autochory and endozoochory)**

Myrmecochory: **non-myrmecochorous (b)**



## Belowground organs and clonality

Shoot metamorphosis: **shoot thorn**

Root metamorphosis: **root shoot**

Position of root buds: **lateral roots**

Role of root buds in life-history of a plant: **additive**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **0**

Number of buds per shoot at the soil surface (root buds included): **0**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**

Size of the belowground bud bank (root buds included): **30**

Depth of the belowground bud bank (root buds included) [cm]: **10**



## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **24**

Ploidy level (x): **2**

2C genome size [Mbp]: **866.01**

1Cx monoploid genome size [Mbp]: **433.01**

Genomic GC content: **38.5 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7x - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area (generalist)**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.72**

Herb layer disturbance frequency indicator value: **-1.05**

Whole-community disturbance severity indicator value: **0.24**

Herb layer disturbance severity indicator value: **0.13**

Whole-community structure based disturbance indicator value: **0.17**

Herb layer structure-based disturbance indicator value: **0.26**

## Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1D Mobile calcareous screes: **1 - rare occurrence**

8 Dry grasslands

8B Submediterranean dry grasslands on rock outcrops: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

11 Heathlands and scrub

11J Willow galleries of loamy and sandy river banks: **1 - rare occurrence**

11L Tall mesic and xeric shrub: **2 - optimum**

11N Low xeric scrub: **2 - optimum**

11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**

12 Forests





12B Alluvial forests: **1 - rare occurrence**  
 12C Oak-hornbeam forests: **1 - rare occurrence**  
 12D Ravine forests: **1 - rare occurrence**  
 12F Limestone beech forests: **2 - optimum**  
 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**  
 12I Sub-continental thermophilous oak forests: **2 - optimum**  
 12J Acidophilous thermophilous oak forests: **2 - optimum**  
 12K Acidophilous oak forests: **1 - rare occurrence**  
 12L Boreo-continental pine forests: **1 - rare occurrence**  
 12O Peri-Alpidic pine forests: **1 - rare occurrence**  
 12T Robinia pseudacacia plantations: **1 - rare occurrence**  
 12W Pine and larch plantations: **2 - optimum**  
 13 Anthropogenic vegetation  
 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**  
 Affinity to the forest environment  
 Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**  
 Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**  
 Diagnostic taxon  
 Diagnostic taxon of alliances: [KBB Berberidion vulgaris](#), [LCA Quercion pubescenti-petraeae](#)  
 Diagnostic taxon of associations: [KBB04 Pruno spinosae-Ligustretum vulgaris](#), [KBB05 Rhamno catharticae-Cornetum sanguineae](#), [KBC04 Senecioni fuchsii-Coryletum avellanae](#), [LBB04 Primulo veris-Carpinetum betuli](#)  
 Constant taxon  
 Constant taxon of associations: [KBB05 Rhamno catharticae-Cornetum sanguineae](#)  
 Ecological specialization indices  
 Ecological specialization index for all vegetation types: **4.3**  
 Ecological specialization index for non-forest vegetation: **4**  
 Ecological specialization index for forest vegetation: **4.6**  
 Colonization ability  
 Index of colonization success (ICS): **3**  
 Index of colonization potential (ICP): **3**  
 Optimum successional age [years]: **13**



## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**  
 Floristic region: **Europe, Western Asia**  
 Distribution range extension along the continentality gradient: **6**  
 Elevational belt in the Czech Republic: **lowlands, colline belt**  
 Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 516  
 taxon.data.freq\_in\_quad: 1354  
 Commonness in vegetation plots from the Czech Republic  
 Occurrence frequency in vegetation plots: **0.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **14.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **2.3 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **4 %**

Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **20**

Number of narrow habitats in which the taxon has its optimum: **7**

Number of broad habitats in which the taxon occurs: **5**

Number of broad habitats in which the taxon has its optimum: **2**

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**