


# *Solanum villosum*

## Distribution




### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.25-0.5**

Growth form: **annual herb**

Life form: **therophyte**

Life strategy: **R - ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

## Flower

Flowering period [month]: **July-October**

Flower colour: **white**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **rotate**

Calyx fusion: **synsepalous**  
Inflorescence type: **cincinnus**  
Dicliny: **synoecious**  
Generative reproduction type: **mixed mating**  
Pollination syndrome: **insect-pollination, selfing**



### **Fruit, seed and dispersal**

Fruit type: **fleshy fruit - berry**  
Fruit colour: **yellow, red**  
Reproduction type: **only by seed/spores**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot life span (cyclicality): **monocyclic shoots prevailing**  
Primary root: **present**  
Bud bank  
Number of buds per shoot at the soil surface (root buds excluded): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
Size of the belowground bud bank (root buds excluded): **5**  
Depth of the belowground bud bank (root buds excluded) [cm]: **1**  
Number of buds per shoot at the soil surface (root buds included): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**  
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
Size of the belowground bud bank (root buds included): **5**  
Depth of the belowground bud bank (root buds included) [cm]: **1**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Taxon origin**

Origin in the Czech Republic: **neophyte**  
Invasion status: **casual**  
Geographic origin: **Mediterranean**  
Year of the first record in the wild: **1998**  
Period of introduction: **Recent Past (1950-2000)**  
Introduction pathway: **unintentional - industry, unintentional - anthropogenic**

### **Habitat and sociology**

Occurrence in habitats  
13 Anthropogenic vegetation  
13A Annual vegetation of ruderal habitats: **2 - optimum**

13C Annual vegetation of trampled habitats: **1 - rare occurrence**

13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

### **Distribution and frequency**

Floristic zone: **southern temperate, submeridional, meridional**

Floristic region: **Europe, Near East**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 19

taxon.data.freq\_in\_quad: 23

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **3**

Number of narrow habitats in which the taxon has its optimum: **1**

Number of broad habitats in which the taxon occurs: **1**

Number of broad habitats in which the taxon has its optimum: **1**

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**