


# *Stratiotes aloides*

## Distribution




### Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.1-0.5**

Growth form: **clonal herb**

Life form: **hydrophyte**

Life strategy (Pierce method based on leaf traits): **CR**

Life strategy (Pierce method, C-score): **60.4 %**

Life strategy (Pierce method, S-score): **0 %**

Life strategy (Pierce method, R-score): **39.6 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **rosulate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **scleromorphic, hydromorphic**

## Flower

Flowering period [month]: **June-September**

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **white**  
Flower symmetry: **actinomorphic**  
Perianth type: **calyx and corolla**  
Perianth fusion: **free**  
Calyx fusion: **aposepalous**  
Inflorescence type: **flores solitarii**  
Dicliny: **dioecious**  
Generative reproduction type: **alogamy**  
Pollination syndrome: **insect-pollination**

### **Fruit, seed and dispersal**

Fruit type: **dry fruit - capsule**  
Reproduction type: **mostly vegetatively, rarely by seed/spores**  
Dispersal unit (diaspore): **seed, turion, stolon-born plantlet**  
Dispersal strategy: **Wolffia (mainly hydrochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot metamorphosis: **stolon, shoot tuber, turion**  
Storage organ: **stolon, shoot tuber, turion**  
Type of clonal growth organ: **stolon**  
Freely dispersible organs of clonal growth: **present**  
Shoot life span (cyclicity): **monocyclic shoots prevailing**  
Branching type of stem-derived organs of clonal growth: **monopodial**  
Primary root: **absent**  
Persistence of the clonal growth organ [year]:  
Number of clonal offspring:  
Lateral spreading distance by clonal growth [m]:  
Clonal index: **7**

### **Bud bank**

Number of buds per shoot at the soil surface (root buds excluded): **3**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded):  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded):  
Size of the belowground bud bank (root buds excluded): **3**  
Depth of the belowground bud bank (root buds excluded) [cm]:  
Number of buds per shoot at the soil surface (root buds included): **3**  
Number of buds per shoot at a depth of 0–10 cm (root buds included):  
Number of buds per shoot at a depth greater than 10 cm (root buds included):  
Size of the belowground bud bank (root buds included): **3**  
Depth of the belowground bud bank (root buds included) [cm]:

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **40 (24, 48)**

Ploidy level (x): **4**

2C genome size [Mbp]: **9097.13**

1Cx monoploid genome size [Mbp]: **2274.28**

Genomic GC content: **43.6 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **7 - heat indicator, occurring in relatively warm lowlands**

Moisture indicator value: **11 - aquatic plant rooted under water, but at least temporarily with leaves above the surface, or a plant floating on the water surface**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **6 - transition between values 5 and 7**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value:

Herb layer disturbance frequency indicator value:

Whole-community disturbance severity indicator value: **0.2**

Herb layer disturbance severity indicator value: **0.23**

Whole-community structure based disturbance indicator value: **0.56**

Herb layer structure-based disturbance indicator value: **0.59**

## Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3A Macrophytic vegetation of eutrophic and mesotrophic still waters: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4C Eutrophic vegetation of muddy substrata: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of associations: [\*\*VAC02 \*Stratiotetum aloidis\*\*\*](#)

## Constant taxon

Constant taxon of associations: [VAC02 \*Stratiotetum aloidis\*](#)

## Dominant taxon

Dominant taxon of associations: [VAC02 \*Stratiotetum aloidis\*](#)

## Ecological specialization indices

Ecological specialization index for all vegetation types: **6.5**

Ecological specialization index for non-forest vegetation: **6.5**

## Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Western Siberia**

Continentality degree: **6**

Distribution range extension along the continentality gradient: **6**

Elevational belt in the Czech Republic: **lowlands, colline belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **74**

taxon.data.freq\_in\_quad: **93**

## Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**

Occurrence frequency in vegetation plots with a cover above 5%: **63 %**

Occurrence frequency in vegetation plots with a cover above 25%: **40.7 %**

Occurrence frequency in vegetation plots with a cover above 50%: **25.9 %**

Mean percentage cover in vegetation plots: **27.8 %**

Maximum percentage cover in vegetation plots: **88 %**

## Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **2**

Number of narrow habitats in which the taxon has its optimum: **0**

Number of broad habitats in which the taxon occurs: **2**

Number of broad habitats in which the taxon has its optimum: **0**

## Threats and protection

Red List 2017 (national categories): **C1t - critically threatened taxon, declining**

Red List 2017 (IUCN categories): **CR - critically endangered**

Legal protection: **endangered taxon**