


# *Utricularia minor*

## Distribution



Map info	
<span style="color: black;">●</span>	revised records
<span style="color: grey;">●</span>	unrevised records
On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.	



## Habitus and growth type

Height [m]: **0.05-0.5**

Growth form: **clonal herb**

Life form: **hydrophyte**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - forked**

Stipules: **absent**

Petiole: **both present and absent**

Leaf life span: **summer green**

Leaf anatomy: **hydromorphic**

## Flower

Flowering period [month]: **June-September**

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**

Flower colour: **yellow**

Flower symmetry: **zygomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**



Shape of the sympetalous corolla or syntepalous perianth: **bilabiate**  
Calyx fusion: **synsepalous**  
Inflorescence type: **racemus**  
Dicliny: **synoecious**  
Generative reproduction type: **mixed mating**  
Pollination syndrome: **insect-pollination**



### **Fruit, seed and dispersal**

Fruit type: **dry fruit - capsule**  
Reproduction type: **mostly vegetatively, rarely by seed/spores**  
Dispersal unit (diaspore): **seed, turion, shoot fragment**  
Dispersal strategy: **Wolffia (mainly hydrochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot metamorphosis: **turion**  
Storage organ: **turion**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Karyology**

Chromosome number (2n): **44 (36, 40)**  
Ploidy level (x): **2**  
2C genome size [Mbp]: **380.23**  
1Cx monoploid genome size [Mbp]: **190.12**  
Genomic GC content: **38.8 %**

### **Taxon origin**

Origin in the Czech Republic: **native**

### **Ecological indicator values**

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **12 - permanently or almost permanently submerged aquatic plant**

Reaction indicator value: **5 - indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions**

Nutrient indicator value: **2 - transition between values 1 and 3**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

## Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **2 - optimum**

5 Vegetation of springs and mires

5E Acidic moss-rich fens and peatland meadows: **2 - optimum**

5F Transitional mires: **2 - optimum**

5H Wet peat soils and bog hollows: **2 - optimum**

11 Heathlands and scrub

11I Willow carrs: **1 - rare occurrence**

12 Forests

12A Alder carrs: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of alliances: [VDC \*Sphagno-Utricularion\*](#)

Diagnostic taxon of associations: [MCG03 \*Peucedano palustris-Caricetum lasiocarpae\*](#), [RBA04 \*Campylio stellati-Caricetum lasiocarpae\*](#), [RBB02 \*Campylio stellati-Trichophoretum alpini\*](#), [RBB03 \*Menyantho trifoliatae-Sphagnetum teretis\*](#), [RBC02 \*Drosero anglicae-Rhynchosporium albae\*](#), [VDC03 \*Scorpidio scorpioidis-Utricularietum\*](#)

Constant taxon

Constant taxon of associations: [VDC03 \*Scorpidio scorpioidis-Utricularietum\*](#)

Dominant taxon

Dominant taxon of associations: [MCG05 \*Caricetum diandrae\*](#)

## Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional, meridional**

Floristic region: **circumpolar**

Distribution range extension along the continentality gradient: **8**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt (montane belt)**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **90**

taxon.data.freq\_in\_quad: 131

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**

Number of narrow habitats in which the taxon has its optimum: **4**

Number of broad habitats in which the taxon occurs: **4**

Number of broad habitats in which the taxon has its optimum: **2**

## **Threats and protection**

Red List 2017 (national categories): **C2b - endangered taxon, rare and declining**

Red List 2017 (IUCN categories): **VU - vulnerable**

Legal protection: **not protected by law**