


# *Cytisus nigricans* subsp. *nigricans*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.3**

Growth form: **shrub**

Life form: **nanophanerophyte**

Life strategy: **C - competitor**

Life strategy (Pierce method based on leaf traits): **S/CSR**

Life strategy (Pierce method, C-score): **11.7 %**

Life strategy (Pierce method, S-score): **58 %**

Life strategy (Pierce method, R-score): **30.3 %**



## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - ternate**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf deciduousness in woody plants: **winter deciduous**

Leaf anatomy: **mesomorphic**

Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

## Flower

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**

Flower colour: **yellow**

Flower symmetry: **zygomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **synsepalous**

Inflorescence type: **racemus**

Dicliny: **synoecious**

Generative reproduction type: **allogamy self-incompatibility**

Pollination syndrome: **insect-pollination**



## Fruit, seed and dispersal

Fruit type: **dry fruit - legume**

Fruit colour: **brown, black**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **myrmecochorous**

## Belowground organs and clonality

Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded):

Number of buds per shoot at a depth of 0–10 cm (root buds excluded):

Number of buds per shoot at a depth greater than 10 cm (root buds excluded):

Size of the belowground bud bank (root buds excluded):

Depth of the belowground bud bank (root buds excluded) [cm]:

Number of buds per shoot at the soil surface (root buds included):

Number of buds per shoot at a depth of 0–10 cm (root buds included):

Number of buds per shoot at a depth greater than 10 cm (root buds included):

Size of the belowground bud bank (root buds included):

Depth of the belowground bud bank (root buds included) [cm]:

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Karyology

Chromosome number (2n): **48**

Ploidy level (x): **4**

2C genome size [Mbp]: **1755.99**

1Cx monoploid genome size [Mbp]: **439**

Genomic GC content: **37.6 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **6 - transition between values 5 and 7**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.71**

Herb layer disturbance frequency indicator value: **-0.71**

Whole-community disturbance severity indicator value: **0.23**

Herb layer disturbance severity indicator value: **0.15**

Whole-community structure based disturbance indicator value: **0.23**

Herb layer structure-based disturbance indicator value: **0.29**

## Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 - rare occurrence**

1B Siliceous cliffs and block fields: **1 - rare occurrence**

1D Mobile calcareous screes: **1 - rare occurrence**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 - rare occurrence**

7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 - rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**

8B Submediterranean dry grasslands on rock outcrops: **2 - optimum**

8C Narrow-leaved sub-continental steppes: **1 - rare occurrence**

8D Broad-leaved dry grasslands: **1 - rare occurrence**

8E Acidophilous dry grasslands: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **2 - optimum**

9 Sand grasslands and rock-outcrop vegetation

9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**

9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**

11 Heathlands and scrub



11A Dry lowland to subalpine heathlands: **2 - optimum**  
 11L Tall mesic and xeric shrub: **1 - rare occurrence**  
 11N Low xeric scrub: **2 - optimum**  
 12 Forests  
 12C Oak-hornbeam forests: **1 - rare occurrence**  
 12D Ravine forests: **1 - rare occurrence**  
 12F Limestone beech forests: **1 - rare occurrence**  
 12G Acidophilous beech forests: **1 - rare occurrence**  
 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**  
 12I Sub-continental thermophilous oak forests: **2 - optimum**  
 12J Acidophilous thermophilous oak forests: **2 - optimum**  
 12K Acidophilous oak forests: **2 - optimum**  
 12L Boreo-continental pine forests: **2 - optimum**  
 12O Peri-Alpidic pine forests: **2 - optimum**  
 12V Spruce plantations: **1 - rare occurrence**  
 12W Pine and larch plantations: **2 - optimum**  
 13 Anthropogenic vegetation  
 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**  
 Affinity to the forest environment  
 Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**  
 Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**  
 Diagnostic taxon  
 Diagnostic taxon of classes: [LD Quercetea robori-petraeae](#)  
 Diagnostic taxon of alliances: [LDA Quercion roboris](#)  
 Diagnostic taxon of associations: [LDA02 Viscario vulgaris-Quercetum petraeae](#),  
[THC04 Asplenio cuneifolii-Seslerietum caeruleae](#)  
 Constant taxon  
 Constant taxon of associations: [LDA02 Viscario vulgaris-Quercetum petraeae](#),  
[THC04 Asplenio cuneifolii-Seslerietum caeruleae](#)  
 Ecological specialization indices  
 Ecological specialization index for all vegetation types: **4.2**  
 Ecological specialization index for non-forest vegetation: **4.2**  
 Ecological specialization index for forest vegetation: **4.8**  
 Colonization ability  
 Index of colonization success (ICS): **4**  
 Index of colonization potential (ICP): **4**  
 Optimum successional age [years]: **75**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**  
 Floristic region: **Europe**  
 Continentality degree: **6**  
 Distribution range extension along the continentality gradient: **2**  
 Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt,**

### **montane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 426

taxon.data.freq\_in\_quad: 1106

### **Commonness in vegetation plots from the Czech Republic**

Occurrence frequency in vegetation plots: **0.8 %**

Occurrence frequency in vegetation plots with a cover above 5%: **9.8 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0.9 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0.4 %**

Mean percentage cover in vegetation plots: **3.5 %**

Maximum percentage cover in vegetation plots: **88 %**

### **Number of habitats with taxon occurrence in the Czech Republic**

Number of narrow habitats in which the taxon occurs: **29**

Number of narrow habitats in which the taxon has its optimum: **11**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **3**

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**