


Chenopodium album subsp. *album*

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.1-1.8**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**



Flower

Flowering period [month]: **July-September**

Flower colour: **green**

Flower symmetry: **actinomorphic**

Perianth type: **homochlamydeous**

Perianth fusion: **fused**

Inflorescence type: **panicula e pseudospicis composita, pseudospica**

Generative reproduction type: **mixed mating**



Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**

Fruit colour: **green, yellow, brown**

Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **5x - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **6x - transition between values 5 and 7 (generalist)**

Nutrient indicator value: **7 - occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4B Halophilous reed and sedge beds: **1 - rare occurrence**

4D Riverine reed vegetation: **1 - rare occurrence**

4E Reed vegetation of brooks: **1 - rare occurrence**

4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**

4I Vegetation of nitrophilous annual hygrophilous herbs: **1 - rare occurrence**

4L Nitrophilous herbaceous fringes of lowland rivers: **1 - rare occurrence**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 - rare occurrence**

6G Vegetation of wet disturbed soils: **1 - rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**

8C Narrow-leaved sub-continental steppes: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**



9 Sand grasslands and rock-outcrop vegetation9B Open vegetation of acidic sands: **1 - rare occurrence**9C Festuca grasslands on acidic sands: **1 - rare occurrence****10 Saline vegetation**10G Continental vegetation of annual halophilous grasses: **1 - rare occurrence**10I Inland saline meadows: **1 - rare occurrence****11 Heathlands and scrub**11J Willow galleries of loamy and sandy river banks: **1 - rare occurrence**11L Tall mesic and xeric shrub: **1 - rare occurrence**11N Low xeric scrub: **1 - rare occurrence**11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence****12 Forests**12T Robinia pseudacacia plantations: **1 - rare occurrence**12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**12V Spruce plantations: **1 - rare occurrence**12W Pine and larch plantations: **1 - rare occurrence****13 Anthropogenic vegetation**13A Annual vegetation of ruderal habitats: **4 - constant dominant**13B Annual vegetation of arable land: **2 - optimum**13C Annual vegetation of trampled habitats: **2 - optimum**13D Perennial thermophilous ruderal vegetation: **2 - optimum**13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence****Affinity to the forest environment**Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests****Distribution and frequency**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 556

taxon.data.freq_in_quad: 1389

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **30**Number of narrow habitats in which the taxon has its optimum: **4**Number of broad habitats in which the taxon occurs: **9**Number of broad habitats in which the taxon has its optimum: **1****Threats and protection**Red List 2017 (national categories): **taxon is not on the Red List**Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**Legal protection: **not protected by law**

