×*Tripleurothemis maleolens*

Distribution



Habitus and growth type

Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Petiole: **both present and absent**

Flower

Flower colour: **white, yellow**
Perianth type: **calyx reduced, corolla present**
Calyx fusion: **pappus**
Inflorescence type: **corymbus ex anthodiis compositus**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **archaeophyte**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Invasion status: **casual**
Geographic origin: **hybrid origin**
Year of the first record in the wild: **1886**
Year of the last record: **1886**
Period of introduction: **Neolithic (5600–4200 BCE)**
Introduction pathway: **unintentional – anthropogenic**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz