


Fraxinus americana agg.

Distribution



© Dana Holubová

Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.


Habitus and growth type

Height [m]: **15-25**

Growth form: **tree**

Life form: **macrophanerophyte**

Life strategy: **C - competitor**



© Dana Holubová

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **opposite**

Leaf shape: **compound - imparipinnate**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf deciduousness in woody plants: **winter deciduous**

Leaf anatomy: **mesomorphic**

Functional leaf type in woody plants: **broad deciduous or semi-deciduous**



© Dana Michalčová

Flower

Flowering period [month]: **April**

Flower colour: **green-white**

Perianth type: **flower achlamydeous**



Calyx fusion: **synsepalous**
 Inflorescence type: **panicula**
 Dicliny: **dioecious**
 Generative reproduction type: **allogamy**
 Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**
 Reproduction type: **mostly by seed/spores, rarely vegetatively**
 Dispersal unit (diaspore): **fruit, infrutescence or its part**
 Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
 Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Root metamorphosis: **root shoot**
 Bud bank
 Number of buds per shoot at the soil surface (root buds excluded): **15**
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
 Size of the belowground bud bank (root buds excluded): **15**
 Depth of the belowground bud bank (root buds excluded) [cm]: **1**
 Number of buds per shoot at the soil surface (root buds included): **15**
 Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
 Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
 Size of the belowground bud bank (root buds included): **15**
 Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode


Parasitism and mycoheterotrophy: **autotrophic**
 Carnivory: **non-carnivorous**
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
 Invasion status: **invasive**
 Geographic origin: **North America**

Habitat and sociology

Occurrence in habitats
 11 Heathlands and scrub
 11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**
 12 Forests
 12B Alluvial forests: **2 - optimum**
 12U Plantations of broad-leaved non-native trees: **3 - dominant**



13 Anthropogenic vegetation

13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Eastern America**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 63

taxon.data.freq_in_quad: 82

