


Vaccinium oxycoccos agg.

Distribution




© Vladimír Nejšpílek

Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Barbora Olšáková

Habitus and growth type

Height [m]: **0.01-0.06**

Growth form: **dwarf shrub**

Life form: **chamaephyte**

Life strategy: **CS - competitor/stress-tolerator**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**

Leaf life span: **evergreen**

Leaf deciduousness in woody plants: **evergreen**

Leaf anatomy: **scleromorphic**

Functional leaf type in woody plants: **sclerophyllous**

Flower

Flowering period [month]: **June-July**

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**

Flower colour: **pink**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **rotate**

Calyx fusion: **synsepalous**

Inflorescence type: **racemus, flores solitarii**

Dicliny: **synoecious**

Generative reproduction type: **mixed mating**

Pollination syndrome: **wind-pollination, insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **fleshy fruit - berry**

Fruit colour: **red**

Reproduction type: **mostly vegetatively, rarely by seed/spores**

Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**

Dispersal strategy: **Cornus (mainly autochory and endozoochory)**

Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon**

Type of clonal growth organ: **stolon, epigeogenous rhizome**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **absent**

Persistence of the clonal growth organ [year]: **4**

Number of clonal offspring: **0.5**

Lateral spreading distance by clonal growth [m]: **0.13**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **15**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **30**

Depth of the belowground bud bank (root buds excluded) [cm]: **3**

Number of buds per shoot at the soil surface (root buds included): **15**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **30**

Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **24, 48, 60, 72**

Ploidy level (x): **2, 4, 5, 6**

2C genome size [Mbp]: **2842.98**

1Cx monoploid genome size [Mbp]: **642.78**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7x - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area (generalist)**

Temperature indicator value: **4 - transition between values 3 and 5**

Moisture indicator value: **9 - wetness indicator, focus on often soaked, poorly aerated soils**

Reaction indicator value: **2x - transition between values 1 and 3 (generalist)**

Nutrient indicator value: **1 - occurring at nutrient-poorest sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.8**

Herb layer disturbance frequency indicator value: **-1.65**

Whole-community disturbance severity indicator value: **0.04**

Herb layer disturbance severity indicator value: **0.05**

Whole-community structure based disturbance indicator value: **0.42**

Herb layer structure-based disturbance indicator value: **0.33**

Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4G Tall-sedge beds: **1 - rare occurrence**

5 Vegetation of springs and mires

5E Acidic moss-rich fens and peatland meadows: **1 - rare occurrence**

5F Transitional mires: **2 - optimum**

5G Raised bogs: **2 - optimum**

5H Wet peat soils and bog hollows: **2 - optimum**

6 Meadows and mesic pastures

6F Intermittently wet Molinia meadows: **1 - rare occurrence**

12 Forests

12P Peatland pine forests: **2 - optimum**

12Q Peatland birch forests: **2 - optimum**



12R Acidophilous spruce forests: **2 - optimum**

Diagnostic taxon

Diagnostic taxon of classes: [RC *Oxycocco-Sphagnetea*](#)

Diagnostic taxon of alliances: [RBD *Sphagno-Caricion canescentis*](#), [RBE *Sphagnion cuspidati*](#), [RCA *Sphagnion magellanicum*](#), [RCB *Oxycocco palustris-Ericion tetralicis*](#), [RCC *Oxycocco microcarpi-Empetrium hermaphroditi*](#)

Diagnostic taxon of associations: [LFD01 *Vaccinio uliginosi-Betuletum pubescentis*](#), [RBC02 *Drosero anglicae-Rhynchosporium albae*](#), [RBD02 *Sphagno recurvi-Caricetum lasiocarpae*](#), [RBE01 *Drepanoclado fluitantis-Caricetum limosae*](#), [RBE03 *Rhynchosporium albae-Sphagnetum tenelli*](#), [RCA01 *Eriophoro vaginati-Sphagnetum recurvi*](#), [RCA02 *Andromeda polifoliae-Sphagnetum magellanicum*](#), [RCA03 *Vaccinio uliginosi-Pinetum mugo*](#), [RCA05 *Ledo palustris-Pinetum uncinatae*](#), [RCB01 *Trichophoro cespitosi-Sphagnetum papillosum*](#), [RCC02 *Empetrum nigri-Sphagnetum fuscum*](#)

Constant taxon

Constant taxon of classes: [RC *Oxycocco-Sphagnetea*](#)

Constant taxon of alliances: [RBE *Sphagnion cuspidati*](#), [RCA *Sphagnion magellanicum*](#), [RCB *Oxycocco palustris-Ericion tetralicis*](#), [RCC *Oxycocco microcarpi-Empetrium hermaphroditi*](#)

Constant taxon of associations: [LFD01 *Vaccinio uliginosi-Betuletum pubescentis*](#), [RBD02 *Sphagno recurvi-Caricetum lasiocarpae*](#), [RBE01 *Drepanoclado fluitantis-Caricetum limosae*](#), [RBE03 *Rhynchosporium albae-Sphagnetum tenelli*](#), [RCA01 *Eriophoro vaginati-Sphagnetum recurvi*](#), [RCA02 *Andromeda polifoliae-Sphagnetum magellanicum*](#), [RCA03 *Vaccinio uliginosi-Pinetum mugo*](#), [RCA04 *Sphagno-Pinetum sylvestris*](#), [RCA05 *Ledo palustris-Pinetum uncinatae*](#), [RCB01 *Trichophoro cespitosi-Sphagnetum papillosum*](#), [RCC02 *Empetrum nigri-Sphagnetum fuscum*](#)

Dominant taxon

Dominant taxon of associations: [MCG03 *Peucedano palustris-Caricetum lasiocarpae*](#), [RBD02 *Sphagno recurvi-Caricetum lasiocarpae*](#), [RBD04 *Polytricho communis-Molinietum caeruleae*](#), [RCA01 *Eriophoro vaginati-Sphagnetum recurvi*](#), [RCA04 *Sphagno-Pinetum sylvestris*](#), [RCA05 *Ledo palustris-Pinetum uncinatae*](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: 5

Ecological specialization index for non-forest vegetation: 5

Ecological specialization index for forest vegetation: **6.6**

Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional**

Floristic region: **circumpolar**

Elevational belt in the Czech Republic: **colline belt, submontane belt, montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 174

taxon.data.freq_in_quad: 357

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **1.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **32.5 %**

Occurrence frequency in vegetation plots with a cover above 25%: **7 %**

Occurrence frequency in vegetation plots with a cover above 50%: **1.4 %**

Mean percentage cover in vegetation plots: **7.9 %**

Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **10**

Number of narrow habitats in which the taxon has its optimum: **6**

Number of broad habitats in which the taxon occurs: **5**

Number of broad habitats in which the taxon has its optimum: **2**