
Allium flavum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.2–0.6**
Growth form: **clonal herb**
Life form: **geophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf anatomy: **succulent, mesomorphic**

Flower

Flowering period [month]: **July–September**
Flower colour: **yellow**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**
Inflorescence type: **pseudumbella**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Dicliny: **synoecious**
Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (a) nv**

Belowground organs and clonality



Shoot metamorphosis: **bulb**
Storage organ: **bulb**
Type of clonal growth organ: **bulb**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **1**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **5**
Number of buds per shoot at the soil surface (root buds included): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **5**





Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **16**
Ploidy level (x): **2**
2C genome size [Mbp]: **23938.34**
1Cx monoploid genome size [Mbp]: **11969.17**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Genomic GC content: **41.2 %**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**



Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**

Moisture indicator value: **2 – transition between values 1 and 3**


Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **2 – transition between values 1 and 3**

Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.18**



Herb layer disturbance frequency indicator value: **-0.67**

Whole-community disturbance severity indicator value: **0.24**

Herb layer disturbance severity indicator value: **0.25**

Whole-community structure based disturbance indicator value: **0.41**

Herb layer structure-based disturbance indicator value: **0.51**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**

1B Siliceous cliffs and block fields: **1 – rare occurrence**

1D Mobile calcareous screes: **1 – rare occurrence**



8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**

8B Submediterranean dry grasslands on rock outcrops: **2 – optimum**

8C Narrow-leaved sub-continental steppes: **1 – rare occurrence**

8D Broad-leaved dry grasslands: **1 – rare occurrence**

8E Acidophilous dry grasslands: **1 – rare occurrence**

8F Thermophilous forest fringe vegetation: **1 – rare occurrence**



9 Sand grasslands and rock-outcrop vegetation

9E Acidophilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

9F Basiphilous vegetation of spring therophytes and succulents: **2 – optimum**



11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 – rare occurrence**


11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12D Ravine forests: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz





12H Peri-Alpidic basiphilous thermophilous oak forests: **2 – optimum**
12J Acidophilous thermophilous oak forests: **2 – optimum**
12O Peri-Alpidic pine forests: **1 – rare occurrence**
12T Robinia pseudacacia plantations: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of alliances: **SAB *Asplenion cuneifolii***, **TFF *Alysso alyssoidis-Sedion***, **THB *Bromo pannonici-Festucion pallentis***
Diagnostic taxon of associations: **LCC02 *Genisto pilosae-Quercetum petraeae***, **SAB02 *Notholaeno marantae-Sempervivetum hirti***, **SCA03 *Teucrio botryos-Melicetum ciliatae***, **TFF01 *Cerastietum***, **THB01 *Poo badensis-Festucetum pallentis***, **THC02 *Minuartio setaceae-Seslerietum caeruleae***, **THC04 *Asplenio cuneifolii-Seslerietum caeruleae***

Constant taxon

Constant taxon of alliances: **THB *Bromo pannonici-Festucion pallentis***
Constant taxon of associations: **SAB02 *Notholaeno marantae-Sempervivetum hirti***, **SCA03 *Teucrio botryos-Melicetum ciliatae***, **THB01 *Poo badensis-Festucetum pallentis***, **THC02 *Minuartio setaceae-Seslerietum caeruleae***, **THC04 *Asplenio cuneifolii-Seslerietum caeruleae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.9**
Ecological specialization index for non-forest vegetation: **5**
Ecological specialization index for forest vegetation: **5.1**

Colonization ability

Index of colonization success (ICS): **2**
Index of colonization potential (ICP): **2**
Optimum successional age [years]: **18**

Distribution and frequency

Floristic zone: **submeridional, meridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 46
taxon.data.freq_in_quad: 84

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.3 %**
Occurrence frequency in vegetation plots with a cover above 5%: **1.1 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.1 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.5 %**
Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **19**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Number of narrow habitats in which the taxon has its optimum: **5**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **NT – near threatened**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz