



Arabidopsis arenosa

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.15-0.4**

Growth form: **monocarpic perennial non-clonal herb**

Life form: **therophyte (hemicryptophyte)**


Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **R/SR**

Life strategy (Pierce method, C-score): **10.4 %**

Life strategy (Pierce method, S-score): **22.9 %**

Life strategy (Pierce method, R-score): **66.7 %**



Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, rosulate**

Leaf shape: **simple - pinnately divided**

Stipules: **absent**

Petiole: **both present and absent**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic**



Flower

Flowering period [month]: **April-July**

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **white, pink**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **aposepalous**

Inflorescence type: **racemus**

Dicliny: **synoecious**

Generative reproduction type: **allogamy self-incompatibility**

Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit - siliqua**

Fruit colour: **brown**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**

Storage organ: **pleiocorm**

Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**

Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded):

Number of buds per shoot at a depth of 0-10 cm (root buds excluded):

Number of buds per shoot at a depth greater than 10 cm (root buds excluded):

Size of the belowground bud bank (root buds excluded):

Depth of the belowground bud bank (root buds excluded) [cm]:


Number of buds per shoot at the soil surface (root buds included):

Number of buds per shoot at a depth of 0-10 cm (root buds included):

Number of buds per shoot at a depth greater than 10 cm (root buds included):

Size of the belowground bud bank (root buds included):

Depth of the belowground bud bank (root buds included) [cm]:




Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **32**

Ploidy level (x): **4**

2C genome size [Mbp]: **768.67**

1Cx monoploid genome size [Mbp]: **192.17**

Genomic GC content: **39.6 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **6 - transition between values 5 and 7**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.74**

Herb layer disturbance frequency indicator value: **-0.91**

Whole-community disturbance severity indicator value: **0.26**

Herb layer disturbance severity indicator value: **0.16**

Whole-community structure based disturbance indicator value: **0.18**

Herb layer structure-based disturbance indicator value: **0.17**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **2 - optimum**

1B Siliceous cliffs and block fields: **2 - optimum**

1C Walls: **1 - rare occurrence**

1D Mobile calcareous screes: **2 - optimum**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 - optimum**

8B Submediterranean dry grasslands on rock outcrops: **2 - optimum**

8E Acidophilous dry grasslands: **1 - rare occurrence**

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 - rare occurrence**

11L Tall mesic and xeric shrub: **1 - rare occurrence**


11N Low xeric scrub: **1 - rare occurrence**

11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**

12 Forests

12C Oak-hornbeam forests: **1 - rare occurrence**



12D Ravine forests: **2 - optimum**



12E Herb-rich beech forests: **1 - rare occurrence**
 12F Limestone beech forests: **2 - optimum**
 12G Acidophilous beech forests: **1 - rare occurrence**
 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**
 12J Acidophilous thermophilous oak forests: **2 - optimum**
 12K Acidophilous oak forests: **2 - optimum**
 12L Boreo-continental pine forests: **2 - optimum**
 12O Peri-Alpidic pine forests: **2 - optimum**
 12T Robinia pseudacacia plantations: **1 - rare occurrence**
 12V Spruce plantations: **1 - rare occurrence**
 12W Pine and larch plantations: **1 - rare occurrence**
 13 Anthropogenic vegetation
 13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**
 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**
 Diagnostic taxon
 Diagnostic taxon of classes: [SC *Thlaspietea rotundifolii*](#)
 Diagnostic taxon of alliances: [KBG *Euphorbio cyparissiae-Robinion pseudoacaciae*](#),
[SCA *Stipion calamagrostis*](#)
 Diagnostic taxon of associations: [KBG01 *Melico transsilvanicae-Robinetum pseudoacaciae*](#),
[LBF04 *Seslerio albicantis-Tilietum cordatae*](#), [THC03 *Saxifrago paniculatae-Seslerietum caeruleae*](#)
 Constant taxon
 Constant taxon of associations: [KAB03 *Salici purpureae-Myricarietum germanicae*](#)
 Ecological specialization indices
 Ecological specialization index for all vegetation types: **4.2**
 Ecological specialization index for non-forest vegetation: **3.5**
 Ecological specialization index for forest vegetation: **4.9**
 Colonization ability
 Index of colonization success (ICS): **5**
 Index of colonization potential (ICP): **6**
 Optimum successional age [years]: **30**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
 Floristic region: **Europe**
 Continentality degree: **5**
 Distribution range extension along the continentality gradient: **3**
 Elevational belt in the Czech Republic: **colline belt, submontane belt**
 Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **308**
 taxon.data.freq_in_quad: 545
 Commonness in vegetation plots from the Czech Republic
 Occurrence frequency in vegetation plots: **0.8 %**
 Occurrence frequency in vegetation plots with a cover above 5%: **1.8 %**
 Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
 Occurrence frequency in vegetation plots with a cover above 50%: **0 %**



Mean percentage cover in vegetation plots: **2.2 %**

Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **27**

Number of narrow habitats in which the taxon has its optimum: **12**

Number of broad habitats in which the taxon occurs: **5**

Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Legal protection: **not protected by law**

