


# Aster amellus

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.2-0.6**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CS - competitor/stress-tolerator**

Life strategy (Pierce method based on leaf traits): **CSR**

Life strategy (Pierce method, C-score): **31.4 %**

Life strategy (Pierce method, S-score): **35.4 %**

Life strategy (Pierce method, R-score): **33.2 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**


Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **both present and absent**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**



## Flower

Flowering period [month]: **July-October**

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**  
 Flower colour: **white, yellow, blue, blue-violet**  
 Flower symmetry: **actinomorphic, zygomorphic**  
 Perianth type: **calyx reduced, corolla present**  
 Perianth fusion: **fused**  
 Shape of the sympetalous corolla or syntepalous perianth: **ligulate, tubular**  
 Calyx fusion: **pappus**  
 Inflorescence type: **corymbothsus ex anthodiis compositus**  
 Dicliny: **gynomonoecious**  
 Generative reproduction type: **facultative allogamy**  
 Pollination syndrome: **insect-pollination, selfing**



## Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**  
 Fruit colour: **brown**  
 Reproduction type: **by seed/spores and vegetatively**  
 Dispersal unit (diaspore): **fruit, infrutescence or its part**  
 Dispersal strategy: **Epilobium (mainly anemochory and autochory)**  
 Myrmecochory: **probably non-myrmecochorous nv, non-myrmecochorous (b)**



## Belowground organs and clonality

Shoot metamorphosis: **rhizome**  
 Storage organ: **rhizome**  
 Type of clonal growth organ: **hypogeogenous rhizome**  
 Freely dispersible organs of clonal growth: **absent**  
 Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**  
 Branching type of stem-derived organs of clonal growth: **sympodial**  
 Primary root: **absent**  
 Persistence of the clonal growth organ [year]: **4**  
 Number of clonal offspring: **1**  
 Lateral spreading distance by clonal growth [m]: **0.13**  
 Clonal index: **4**  
**Bud bank**  
 Number of buds per shoot at the soil surface (root buds excluded): **5**  
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**  
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
 Size of the belowground bud bank (root buds excluded): **20**  
 Depth of the belowground bud bank (root buds excluded) [cm]: **4**  
 Number of buds per shoot at the soil surface (root buds included): **5**  
 Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**  
 Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
 Size of the belowground bud bank (root buds included): **20**  
 Depth of the belowground bud bank (root buds included) [cm]: **4**



## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **3 - missing on damp soil**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.08**

Herb layer disturbance frequency indicator value: **-0.29**

Whole-community disturbance severity indicator value: **0.2**

Herb layer disturbance severity indicator value: **0.21**

Whole-community structure based disturbance indicator value: **0.47**

Herb layer structure-based disturbance indicator value: **0.5**

## Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1D Mobile calcareous screes: **1 - rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**

8B Submediterranean dry grasslands on rock outcrops: **1 - rare occurrence**

8C Narrow-leaved sub-continental steppes: **2 - optimum**

8D Broad-leaved dry grasslands: **2 - optimum**

8F Thermophilous forest fringe vegetation: **2 - optimum**

9 Sand grasslands and rock-outcrop vegetation

9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 - rare occurrence**

11N Low xeric scrub: **1 - rare occurrence**

12 Forests

12C Oak-hornbeam forests: **1 - rare occurrence**

12D Ravine forests: **1 - rare occurrence**

12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**

12J Acidophilous thermophilous oak forests: **1 - rare occurrence**





12L Boreo-continental pine forests: **1 - rare occurrence**

12O Peri-Alpidic pine forests: **2 - optimum**

12W Pine and larch plantations: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Diagnostic taxon

Diagnostic taxon of alliances: [LCA Quercion pubescenti-petraeae](#), [THE Cirsio-Brachypodium pinnati](#)

Diagnostic taxon of associations: [LCA02 Lithospermo purpureocaerulei-Quercetum pubescentis](#), [THE03 Polygalo majoris-Brachypodietum pinnati](#)

Constant taxon

Constant taxon of associations: [LCA02 Lithospermo purpureocaerulei-Quercetum pubescentis](#), [THE03 Polygalo majoris-Brachypodietum pinnati](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.7**

Ecological specialization index for non-forest vegetation: **5.8**

Ecological specialization index for forest vegetation: **5.5**

Colonization ability

Index of colonization success (ICS): **4**

Index of colonization potential (ICP): **3**

Optimum successional age [years]: **30**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Western Siberia**

Continental degree: **7**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **lowlands, colline belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **114**

taxon.data.freq\_in\_quad: **222**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.5 %**

Occurrence frequency in vegetation plots with a cover above 5%: **11.8 %**

Occurrence frequency in vegetation plots with a cover above 25%: **1.3 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0.7 %**

Mean percentage cover in vegetation plots: **3.8 %**

Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **16**

Number of narrow habitats in which the taxon has its optimum: **5**

Number of broad habitats in which the taxon occurs: **5**

Number of broad habitats in which the taxon has its optimum: **2**



## Threats and protection

Red List 2017 (national categories): **C3 - vulnerable taxon**

Red List 2017 (IUCN categories): **NT - near threatened**

Legal protection: **vulnerable taxon**

