Carex vesicaria

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.3–1.2**
Growth form: **clonal herb**
Life form: **hemicryptophyte (geophyte)**
Life strategy: **CS – competitor/stress-tolerator**
Life strategy (Pierce method based on leaf traits): **S/CSR**
Life strategy (Pierce method, C-score): **26.2 %**
Life strategy (Pierce method, S-score): **53.4 %**
Life strategy (Pierce method, R-score): **20.5 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, helomorphic**

Flower

Flowering period [month]: **May–June**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **4 Fagus sylvatica-Galeobdolon (start of mid-spring)**
Flower colour: **yellow-green**
Perianth type: **flower achlamydeous**
Inflorescence type: **spica e spiculis composita**
Dicliny: **monoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal



Fruit type: **dry fruit – nut enclosed in an utricle**
Fruit colour: **yellow, brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Sparganium (mainly autochory and hydrochory)**
Myrmecochory: **probably non-myrmecochorous**

Belowground organs and clonality

Shoot metamorphosis: **stolon, rhizome**
Storage organ: **stolon, rhizome**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **4.5**
Lateral spreading distance by clonal growth [m]: **0.1**
Clonal index: **5**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Karyology



Chromosome number (2n): **80**
Ploidy level (x): **2**
2C genome size [Mbp]: **772.36**
1Cx monoploid genome size [Mbp]: **386.18**
Genomic GC content: **36.1 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **9 – wetness indicator, focus on often soaked, poorly aerated soils**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.82**
Herb layer disturbance frequency indicator value: **-0.64**
Whole-community disturbance severity indicator value: **0.21**
Herb layer disturbance severity indicator value: **0.24**
Whole-community structure based disturbance indicator value: **0.54**
Herb layer structure-based disturbance indicator value: **0.59**

Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 – rare occurrence**

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 – rare occurrence**
4B Halophilous reed and sedge beds: **1 – rare occurrence**
4C Eutrophic vegetation of muddy substrata: **1 – rare occurrence**
4D Riverine reed vegetation: **1 – rare occurrence**
4E Reed vegetation of brooks: **1 – rare occurrence**
4F Mesotrophic vegetation of muddy substrata: **1 – rare occurrence**
4G Tall-sedge beds: **3 – dominant**
4H Vegetation of low annual hygrophilous herbs: **1 – rare occurrence**
4I Vegetation of nitrophilous annual hygrophilous herbs: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

5 Vegetation of springs and mires

5D Calcareous fens: **1 – rare occurrence**
5E Acidic moss-rich fens and peatland meadows: **2 – optimum**
5F Transitional mires: **1 – rare occurrence**
5H Wet peat soils and bog hollows: **1 – rare occurrence**

6 Meadows and mesic pastures

6D Alluvial meadows of lowland rivers: **1 – rare occurrence**
6E Wet Cirsium meadows: **2 – optimum**
6F Intermittently wet Molinia meadows: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

11 Heathlands and scrub

11I Willow carrs: **2 – optimum**
11J Willow galleries of loamy and sandy river banks: **2 – optimum**

12 Forests

12A Alder carrs: **2 – optimum**
12B Alluvial forests: **1 – rare occurrence**
12Q Peatland birch forests: **1 – rare occurrence**
12R Acidophilous spruce forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of alliances: **MCH *Magno-Caricion gracilis***
Diagnostic taxon of associations: **MCH04 *Caricetum vesicariae***, **TDF13 *Lysimachio vulgaris-Filipenduletum ulmariae***

Constant taxon

Constant taxon of associations: **MCH04 *Caricetum vesicariae***, **TDF13 *Lysimachio vulgaris-Filipenduletum ulmariae***

Dominant taxon

Dominant taxon of associations: **MCH04 *Caricetum vesicariae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.6**
Ecological specialization index for non-forest vegetation: **4.7**
Ecological specialization index for forest vegetation: **5.5**

Colonization ability

Index of colonization success (ICS): **5**
Index of colonization potential (ICP): **3**
Optimum successional age [years]: **11**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **circumpolar**
Distribution range extension along the continentality gradient: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 606

taxon.data.freq_in_quad: 1839

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **2.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **35.1 %**

Occurrence frequency in vegetation plots with a cover above 25%: **22.9 %**

Occurrence frequency in vegetation plots with a cover above 50%: **20.9 %**

Mean percentage cover in vegetation plots: **20 %**

Maximum percentage cover in vegetation plots: **99 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **24**

Number of narrow habitats in which the taxon has its optimum: **6**

Number of broad habitats in which the taxon occurs: **7**

Number of broad habitats in which the taxon has its optimum: **5**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz