


Galium saxatile

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.1-0.3**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CS - competitor/stress-tolerator**

Life strategy (Pierce method based on leaf traits): **R/SR**

Life strategy (Pierce method, C-score): **0 %**

Life strategy (Pierce method, S-score): **23 %**

Life strategy (Pierce method, R-score): **77 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **verticillate**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **absent**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June-July**

Flowering phase: **7 Ligustrum vulgare-Stachys sylvatica (end of early summer)**
Flower colour: **white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx absent, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **rotate**
Inflorescence type: **panicula e dichasiis composita**
Dicliny: **synoecious**
Generative reproduction type: **allogamy self-incompatibility**
Pollination syndrome: **insect-pollination**
Pollinator spectrum: **flies s. l., butterflies (other Hymenoptera, meat flies s. l., other Diptera, beetles)**

Fruit, seed and dispersal

Fruit type: **dry fruit - pair of nutlets**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon, pleiocorm**
Storage organ: **stolon, pleiocorm**
Type of clonal growth organ: **stolon**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.13**
Clonal index: **4**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **10**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds excluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **10**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **10**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **44**

Ploidy level (x): **4**

2C genome size [Mbp]: **2531.77**

1Cx monoploid genome size [Mbp]: **632.94**

Genomic GC content: **41.2 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **4 - transition between values 3 and 5**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **2 - transition between values 1 and 3**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.5**

Herb layer disturbance frequency indicator value: **-0.66**

Whole-community disturbance severity indicator value: **0.31**

Herb layer disturbance severity indicator value: **0.28**

Whole-community structure based disturbance indicator value: **0.25**

Herb layer structure-based disturbance indicator value: **0.35**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1B Siliceous cliffs and block fields: **1 - rare occurrence**

2 Alpine and subalpine grasslands

2A Alpine grasslands on siliceous bedrock: **2 - optimum**

2B Subalpine tall-forb and tall-grass vegetation: **2 - optimum**

5 Vegetation of springs and mires

5C Alpine and subalpine soft-water springs: **1 - rare occurrence**

5F Transitional mires: **1 - rare occurrence**

5G Raised bogs: **1 - rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **2 - optimum**

6C Pastures and park grasslands: **1 - rare occurrence**

6E Wet Cirsium meadows: **1 - rare occurrence**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 - optimum**

7B Submontane Nardus grasslands: **2 - optimum**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **2 - optimum**

11D Subalpine acidophilous Pinus mugo scrub: **1 - rare occurrence**

11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**

12 Forests

12E Herb-rich beech forests: **1 - rare occurrence**

12G Acidophilous beech forests: **1 - rare occurrence**

12K Acidophilous oak forests: **1 - rare occurrence**

12L Boreo-continental pine forests: **2 - optimum**

12R Acidophilous spruce forests: **2 - optimum**

12S Basiphilous spruce forests: **2 - optimum**

12V Spruce plantations: **2 - optimum**

12W Pine and larch plantations: **1 - rare occurrence**

13 Anthropogenic vegetation

13F Herbaceous vegetation of forests clearings and Rubus scrub: **2 - optimum**

Affinity to the forest environment

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Diagnostic taxon

Diagnostic taxon of classes: [AC Elyno-Seslerietea](#)

Diagnostic taxon of alliances: [ABB Nardo strictae-Caricion bigelowii](#), [ACA Agrostion alpinae](#), [ADB Calamagrostion arundinaceae](#), [LFC Piceion abietis](#), [TDB Polygono bistortae-Trisetion flavescentis](#), [TEB Nardo strictae-Agrostion tenuis](#)

Diagnostic taxon of associations: [ABB01 Carici bigelowii-Nardetum strictae](#), [ACA01 Saxifrago oppositifoliae-Festucetum versicoloris](#), [ADB01 Bupleuro longifoliae-Calamagrostietum arundinaceae](#), [TDB03 Meo athamantici-Festucetum rubrae](#), [TEA02 Thesio alpini-Nardetum strictae](#), [TEB01 Sileno vulgaris-Nardetum strictae](#), [XEA02 Digitali purpureae-Epilobietum angustifolii](#), [XEA04 Junco effusi-Calamagrostietum villosae](#)

Constant taxon

Constant taxon of alliances: [TDB Polygono bistortae-Trisetion flavescentis](#), [TEB Nardo strictae-Agrostion tenuis](#)

Constant taxon of associations: [ACA01 Saxifrago oppositifoliae-Festucetum versicoloris](#), [TDB03 Meo athamantici-Festucetum rubrae](#), [TEB01 Sileno vulgaris-Nardetum strictae](#)

Dominant taxon

Dominant taxon of associations: [TDB03 Meo athamantici-Festucetum rubrae](#), [TEB01 Sileno vulgaris-Nardetum strictae](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.8**

Ecological specialization index for non-forest vegetation: **5**

Ecological specialization index for forest vegetation: **6.4**

Colonization ability

Index of colonization success (ICS): **5**

Index of colonization potential (ICP): **5**

Optimum successional age [years]: **8**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe**

Continental degree: **2**

Distribution range extension along the continentality gradient: **2**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt, subalpine belt**

Expansive taxon in the region: **Bohemian Moravian Mesophyticum, Bohemian Moravian Oreophyticum**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **235**

taxon.data.freq_in_quad: **514**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.9 %**

Occurrence frequency in vegetation plots with a cover above 5%: **23.2 %**

Occurrence frequency in vegetation plots with a cover above 25%: **4.2 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **6 %**

Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **23**

Number of narrow habitats in which the taxon has its optimum: **11**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **6**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**