


# *Herniaria glabra*

## Distribution




### Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.01-0.03**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **therophyte (hemicryptophyte)**

Life strategy: **R - ruderal**

Life strategy (Pierce method based on leaf traits): **R/SR**

Life strategy (Pierce method, C-score): **0 %**

Life strategy (Pierce method, S-score): **32.1 %**

Life strategy (Pierce method, R-score): **67.9 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **opposite**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **absent**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **June-September**

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**  
 Flower colour: **green**  
 Perianth type: **calyx present, corolla reduced**  
 Perianth fusion: **reduced**  
 Calyx fusion: **synsepalous**  
 Inflorescence type: **verticillastrum**  
 Dicliny: **synoecious**  
 Generative reproduction type: **facultative allogamy, autogamy**  
 Pollination syndrome: **insect-pollination, selfing**



## Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**  
 Fruit colour: **brown**  
 Reproduction type: **only by seed/spores**  
 Dispersal unit (diaspore): **seed, fruit, infructescence or its part**  
 Dispersal strategy: **Allium (mainly autochory)**  
 Myrmecochory: **probably myrmecochorous**



## Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**  
 Storage organ: **pleiocorm**  
 Shoot life span (cyclicity): **monocyclic shoots prevailing**  
 Branching type of stem-derived organs of clonal growth: **sympodial**  
 Primary root: **present**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded):  
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded):  
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded):  
 Size of the belowground bud bank (root buds excluded):  
 Depth of the belowground bud bank (root buds excluded) [cm]:  
 Number of buds per shoot at the soil surface (root buds included):  
 Number of buds per shoot at a depth of 0–10 cm (root buds included):  
 Number of buds per shoot at a depth greater than 10 cm (root buds included):  
 Size of the belowground bud bank (root buds included):  
 Depth of the belowground bud bank (root buds included) [cm]:

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**  
 Carnivory: **non-carnivorous**  
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **18 (36)**  
 Ploidy level (x): **2 (4)**  
 2C genome size [Mbp]: **1166.49**

1Cx monoploid genome size [Mbp]: **583.24**

Genomic GC content: **41.6 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **9 - full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **3 - missing on damp soil**

Reaction indicator value: **4 - transition between values 3 and 5**

Nutrient indicator value: **4 - transition between values 3 and 5**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **0.32**

Herb layer disturbance frequency indicator value: **0.32**

Whole-community disturbance severity indicator value: **0.66**

Herb layer disturbance severity indicator value: **0.68**

Whole-community structure based disturbance indicator value: **0.76**

Herb layer structure-based disturbance indicator value: **0.74**

## Habitat and sociology

Occurrence in habitats

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 - rare occurrence**

7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 - rare occurrence**

8 Dry grasslands

8E Acidophilous dry grasslands: **1 - rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **2 - optimum**

9C Festuca grasslands on acidic sands: **2 - optimum**

9D Pannonian sand steppes: **2 - optimum**

9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 - rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 - rare occurrence**

13B Annual vegetation of arable land: **1 - rare occurrence**

13C Annual vegetation of trampled habitats: **2 - optimum**

13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: [TG Festucetea vaginatae](#), [XA Polygono arenastri-Poëtea annuae](#)

Diagnostic taxon of alliances: [TFC Armerion elongatae](#), [TGA Festucion vaginatae](#), [XAB Saginion procumbentis](#)

Diagnostic taxon of associations: [TFC01 Sileno otitae-Festucetum brevipilae](#), [TGA01 Diantho serotini-Festucetum vaginatae](#), [XAB02 Herniarietum glabrae](#)

Constant taxon

Constant taxon of associations: [XAB02 Herniarietum glabrae](#)

Dominant taxon

Dominant taxon of associations: [XAB02 Herniarietum glabrae](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.6**

Ecological specialization index for non-forest vegetation: **4.6**

Colonization ability

Index of colonization success (ICS): **2**

Index of colonization potential (ICP): **2**

Optimum successional age [years]: **26**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe, Western Asia**

Continental degree: **7**

Distribution range extension along the continentality gradient: **7**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt (montane belt)**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **341**

taxon.data.freq\_in\_quad: **630**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.3 %**

Occurrence frequency in vegetation plots with a cover above 5%: **44.6 %**

Occurrence frequency in vegetation plots with a cover above 25%: **17.8 %**

Occurrence frequency in vegetation plots with a cover above 50%: **4 %**

Mean percentage cover in vegetation plots: **12.2 %**

Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **12**

Number of narrow habitats in which the taxon has its optimum: **4**

Number of broad habitats in which the taxon occurs: **6**

Number of broad habitats in which the taxon has its optimum: **2**

## **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**