
Lunaria annua

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.3–1**
Growth form: **annual herb**
Life form: **hemicryptophyte**
Life strategy: **CR – competitor/ruderal**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **59.4 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **40.6 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **evergreen**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **April–June**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flower colour: **red-violet**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal



Fruit type: **dry fruit – silicula**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality



Root metamorphosis: **secondary storage root**
Storage organ: **secondary storage root**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **archaeophyte/neophyte**
Invasion status: **naturalized**
Geographic origin: **Mediterranean**
Year of the first record in the wild: **1819**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz


Period of introduction: **Late Middle Ages and Early Modern Period (merged category, 1200–1800)**
Introduction pathway: **intentional – ornamental**



Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **5 – semi-shade plant, only exceptionally occurring in full light, but usually at more than 10% of the diffuse radiation incident in an open area**
Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **7 – occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Habitat and sociology

Occurrence in habitats

11 Heathlands and scrub

11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12B Alluvial forests: **1 – rare occurrence**
12C Oak-hornbeam forests: **1 – rare occurrence**
12D Ravine forests: **1 – rare occurrence**
12T Robinia pseudacacia plantations: **2 – optimum**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**
12V Spruce plantations: **1 – rare occurrence**



13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 – optimum**



Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**



Distribution and frequency

Floristic zone: **submeridional, meridional**
Floristic region: **Europe**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 200
taxon.data.freq_in_quad: 275

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **10**

Number of narrow habitats in which the taxon has its optimum: **2**

Number of broad habitats in which the taxon occurs: **3**

Number of broad habitats in which the taxon has its optimum: **2**



Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**









© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz