


# *Microrrhinum minus*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.05-0.25**

Growth form: **annual herb**

Life form: **therophyte**

Life strategy: **R - ruderal**

Life strategy (Pierce method based on leaf traits): **R**

Life strategy (Pierce method, C-score): **8.7 %**

Life strategy (Pierce method, S-score): **0 %**

Life strategy (Pierce method, R-score): **91.4 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, opposite**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **scleromorphic, mesomorphic**



## Flower

Flowering period [month]: **June-September**

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **violet**

Flower symmetry: **zygomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **personate**


Calyx fusion: **synsepalous**

Inflorescence type: **racemus**

Dicliny: **synoecious**

Generative reproduction type: **autogamy**

Pollination syndrome: **selfing**



## Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**

Fruit colour: **brown**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **seed**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **probably myrmecochorous**

## Belowground organs and clonality

Shoot life span (cyclicality): **monocyclic shoots prevailing**

Primary root: **present**

Position of root buds: **hypocotyl**

Role of root buds in life-history of a plant: **additive**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **2**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded):

Number of buds per shoot at a depth greater than 10 cm (root buds excluded):

Size of the belowground bud bank (root buds excluded): **2**

Depth of the belowground bud bank (root buds excluded) [cm]:

Number of buds per shoot at the soil surface (root buds included): **9**

Number of buds per shoot at a depth of 0–10 cm (root buds included):

Number of buds per shoot at a depth greater than 10 cm (root buds included):

Size of the belowground bud bank (root buds included): **9**

Depth of the belowground bud bank (root buds included) [cm]:



## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **14**

Ploidy level (x): **2**

2C genome size [Mbp]: **784.83**  
 1Cx monoploid genome size [Mbp]: **392.41**  
 Genomic GC content: **38.9 %**

## Taxon origin

Origin in the Czech Republic: **archaeophyte**  
 Invasion status: **naturalized**  
 Geographic origin: **Europe, Mediterranean**  
 Period of introduction: **Early Middle Ages (550-1200)**  
 Introduction pathway: **unintentional - anthropogenic**

## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **9 - full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**  
 Temperature indicator value: **6 - transition between values 5 and 7**  
 Moisture indicator value: **4 - transition between values 3 and 5**  
 Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**  
 Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**  
 Salinity indicator value: **0 - not salt tolerant, glycophyte**  
 Indicator values for disturbance  
 Whole-community disturbance frequency indicator value: **-0.03**  
 Herb layer disturbance frequency indicator value: **-0.02**  
 Whole-community disturbance severity indicator value: **0.91**  
 Herb layer disturbance severity indicator value: **0.91**  
 Whole-community structure based disturbance indicator value: **0.77**  
 Herb layer structure-based disturbance indicator value: **0.85**

## Habitat and sociology

### Occurrence in habitats

1 Vegetation of cliffs, screes and walls  
 1A Calcareous cliffs: **1 - rare occurrence**  
 1D Mobile calcareous screes: **2 - optimum**  
 9 Sand grasslands and rock-outcrop vegetation  
 9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**  
 13 Anthropogenic vegetation  
 13A Annual vegetation of ruderal habitats: **1 - rare occurrence**  
 13B Annual vegetation of arable land: **2 - optimum**  
 13C Annual vegetation of trampled habitats: **2 - optimum**  
 13D Perennial thermophilous ruderal vegetation: **2 - optimum**

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**



Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: [SC \*Thlaspietea rotundifolii\*](#)

Diagnostic taxon of alliances: [SCA \*Stipion calamagrostis\*](#), [XBJ \*Salsolion ruthenicae\*](#)

Diagnostic taxon of associations: [SCA02 \*Galeopsietum angustifoliae\*](#), [XBJ01 \*Chenopodietum botryos\*](#)

Constant taxon

Constant taxon of associations: [KAB03 \*Salici purpureae-Myricarietum germanicae\*](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.1**

Ecological specialization index for non-forest vegetation: **5**

Colonization ability

Index of colonization success (ICS): **6**

Index of colonization potential (ICP): **8**

Optimum successional age [years]: **4**



## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe**

Continental degree: **4**

Distribution range extension along the continentality gradient: **4**

Elevational belt in the Czech Republic: **lowlands, colline belt (montane belt)**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **486**

taxon.data.freq\_in\_quad: **1029**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.5 %**

Occurrence frequency in vegetation plots with a cover above 5%: **1.3 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **2.2 %**

Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **7**

Number of narrow habitats in which the taxon has its optimum: **4**

Number of broad habitats in which the taxon occurs: **3**

Number of broad habitats in which the taxon has its optimum: **2**

## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**