


Panicum dichotomiflorum

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.3-1**

Life form: **therophyte**

Life strategy: **CR - competitor/ruderal**

Life strategy (Pierce method based on leaf traits): **C/CSR**

Life strategy (Pierce method, C-score): **61.3 %**

Life strategy (Pierce method, S-score): **14.6 %**

Life strategy (Pierce method, R-score): **24.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **July-September**

Flower colour: **green**

Perianth type: **reduced**
 Perianth fusion: **reduced**
 Inflorescence type: **panicula e spiculis composita**
 Dicliny: **synoecious**
 Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - caryopsis**
 Reproduction type: **only by seed/spores**
 Dispersal unit (diaspore): **fruit, infrutescence or its part**
 Dispersal strategy: **Allium (mainly autochory)**
 Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
 Carnivory: **non-carnivorous**
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
 Invasion status: **casual**
 Geographic origin: **North America**
 Year of the first record in the wild: **1970**
 Period of introduction: **Recent Past (1950-2000)**
 Introduction pathway: **unintentional - traffic**

Habitat and sociology

Affinity to the forest environment
 Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**
 Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
 Floristic region: **Eastern America**
 Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 2
 taxon.data.freq_in_quad: 2

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
 Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red**



List)

Legal protection: **not protected by law**