


Poa annua

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.05-0.25**

Growth form: **annual herb**

Life form: **hemicryptophyte (therophyte)**

Life strategy: **R - ruderal**

Life strategy (Pierce method based on leaf traits): **SR**

Life strategy (Pierce method, C-score): **6.8 %**

Life strategy (Pierce method, S-score): **35 %**

Life strategy (Pierce method, R-score): **58.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic, hygromorphic**

Flower

Flowering period [month]: **January-December**

Flowering phase: **2 Acer platanoides-Anemone nemorosa (start of early spring)**

Flower colour: **green**

Perianth type: **reduced**

Perianth fusion: **reduced**

Inflorescence type: **panicula e spiculis composita**

Dicliny: **synoecious, gynomonoecious**

Generative reproduction type: **facultative autogamy**

Pollination syndrome: **wind-pollination, selfing, cleistogamy**

Fruit, seed and dispersal

Fruit type: **dry fruit - caryopsis**

Fruit colour: **brown**

Reproduction type: **mostly by seed/spores, rarely vegetatively**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (a)**

Belowground organs and clonality

Shoot metamorphosis: **stolon**

Storage organ: **stolon, tuft**

Shoot life span (cyclicality): **monocyclic shoots prevailing**

Primary root: **absent**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**

Number of buds per shoot at a depth of 0-10 cm (root buds excluded): **3**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **8**

Depth of the belowground bud bank (root buds excluded) [cm]: **3**

Number of buds per shoot at the soil surface (root buds included): **5**

Number of buds per shoot at a depth of 0-10 cm (root buds included): **3**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **8**

Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology



Chromosome number (2n): **28**

Ploidy level (x): **4**

2C genome size [Mbp]: **3785.77**

1Cx monoploid genome size [Mbp]: **946.44**

Genomic GC content: **46.4 %**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **5x - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**

Moisture indicator value: **6 - transition between values 5 and 7**

Reaction indicator value: **6x - transition between values 5 and 7 (generalist)**

Nutrient indicator value: **8 - pronounced nutrient indicator**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.21**

Herb layer disturbance frequency indicator value: **0.01**

Whole-community disturbance severity indicator value: **0.77**

Herb layer disturbance severity indicator value: **0.78**

Whole-community structure based disturbance indicator value: **0.68**

Herb layer structure-based disturbance indicator value: **0.71**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 - rare occurrence**

1B Siliceous cliffs and block fields: **1 - rare occurrence**

1C Walls: **1 - rare occurrence**

1D Mobile calcareous screes: **1 - rare occurrence**

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**

3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4D Riverine reed vegetation: **1 - rare occurrence**

4E Reed vegetation of brooks: **1 - rare occurrence**

4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**

4I Vegetation of nitrophilous annual hygrophilous herbs: **2 - optimum**

4J River gravel banks: **1 - rare occurrence**

4K Petasites fringes of montane brooks: **1 - rare occurrence**

4L Nitrophilous herbaceous fringes of lowland rivers: **1 - rare occurrence**

5 Vegetation of springs and mires

5A Hard-water springs with tufa formation: **1 - rare occurrence**

5B Lowland to montane soft-water springs: **1 - rare occurrence**



6 Meadows and mesic pastures

6C Pastures and park grasslands: **2 - optimum**6G Vegetation of wet disturbed soils: **2 - optimum**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **1 - rare occurrence**7B Submontane *Nardus* grasslands: **1 - rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 - rare occurrence**9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**9F Basiphilous vegetation of spring therophytes and succulents: **1 - rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 - rare occurrence**10J Saline steppes: **1 - rare occurrence**

11 Heathlands and scrub

11J Willow galleries of loamy and sandy river banks: **1 - rare occurrence**11L Tall mesic and xeric shrub: **1 - rare occurrence**11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**

12 Forests

12T Robinia pseudacacia plantations: **1 - rare occurrence**12V Spruce plantations: **1 - rare occurrence**12W Pine and larch plantations: **1 - rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **2 - optimum**13B Annual vegetation of arable land: **2 - optimum**13C Annual vegetation of trampled habitats: **4 - constant dominant**13D Perennial thermophilous ruderal vegetation: **2 - optimum**13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 - optimum**13F Herbaceous vegetation of forests clearings and *Rubus* scrub: **1 - rare occurrence**


Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: [*XA Polygono arenastri-Poëtea annuae*](#)Diagnostic taxon of alliances: [*XAA Coronopodo-Polygonion arenastri*](#), [*XAB Saginion procumbentis*](#), [*XBI Malvion neglectae*](#)Diagnostic taxon of associations: [*TDC04 Prunello vulgaris-Ranunculetum repentis*](#), [*XAB01 Sagino procumbentis-Bryetum argentei*](#), [*XAB04 Poëtum annuae*](#), [*XAB05 Lolio perennis-Matricarietum discoideae*](#), [*XBI05 Matricario discoideae-Anthemidetum cotulae*](#)

Constant taxon

Constant taxon of classes: [*XA Polygono arenastri-Poëtea annuae*](#)Constant taxon of alliances: [*TDC Cynosurion cristati*](#), [*XAA Coronopodo-Polygonion arenastri*](#), [*XAB Saginion procumbentis*](#), [*XBC Scleranthion annui*](#), [*XBI Malvion neglectae*](#)Constant taxon of associations: [*MAC02 Cerastio dubii-Ranunculetum sardoi*](#),

[MBB01 *Chenopodietum rubri*](#), [TDC03 *Lolietum perennis*](#), [TDC04 *Prunello vulgaris-Ranunculetum repentis*](#), [TDC05 *Alchemillo hybridae-Poëtum supinae*](#), [TFB01 *Airetum praecocis*](#), [XAA01 *Polygonetum arenastri*](#), [XAA03 *Poa annuae-Coronopodetum squamati*](#), [XAA04 *Eragrostio minoris-Polygonetum arenastri*](#), [XAB01 *Sagino procumbentis-Bryetum argentei*](#), [XAB02 *Herniarietum glabrae*](#), [XAB03 *Rumici acetosellae-Spergularietum rubrae*](#), [XAB04 *Poëtum annuae*](#), [XAB05 *Lolio perennis-Matricarietum discoideae*](#), [XBC01 *Aphano arvensis-Matricarietum chamomillae*](#), [XBC02 *Spergulo arvensis-Scleranthetum annui*](#), [XBC03 *Erophilo verna-Arabidopsietum thalianae*](#), [XBG02 *Chenopodietum urbici*](#), [XBG06 *Atriplicetum roseae*](#), [XBG10 *Chamaeplietum officinalis*](#), [XBI02 *Malvetum pusillae*](#), [XBI04 *Malvo neglectae-Chenopodietum vulvariae*](#), [XBI05 *Matricario discoideae-Anthemidetum cotulae*](#)

Dominant taxon

Dominant taxon of associations: [TDC04 *Prunello vulgaris-Ranunculetum repentis*](#), [XAB04 *Poëtum annuae*](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.2**

Ecological specialization index for non-forest vegetation: **4.3**

Ecological specialization index for forest vegetation: **3.7**

Colonization ability

Index of colonization success (ICS): **7**

Index of colonization potential (ICP): **4**

Optimum successional age [years]: **4**

Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional, meridional, subtropical, tropical, austral or antarctic**

Floristic region: **Europe, circumpolar**

Distribution range extension along the continentality gradient: **8**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **650**

taxon.data.freq_in_quad: **2161**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **5.8 %**

Occurrence frequency in vegetation plots with a cover above 5%: **22.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **10.5 %**

Occurrence frequency in vegetation plots with a cover above 50%: **4.6 %**

Mean percentage cover in vegetation plots: **9 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **36**

Number of narrow habitats in which the taxon has its optimum: **8**

Number of broad habitats in which the taxon occurs: **12**

Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**