



# Potamogeton pusillus

## Distribution



Map info

- revised records
- unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.1-1**

Growth form: **clonal herb**

Life form: **hydrophyte**

Life strategy (Pierce method based on leaf traits): **SR**

Life strategy (Pierce method, C-score): **0.5 %**

Life strategy (Pierce method, S-score): **42.5 %**

Life strategy (Pierce method, R-score): **57 %**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **mainly absent**

Leaf life span: **summer green**

Leaf anatomy: **hydromorphic**

## Flower

Flowering period [month]: **June-August**

Flower colour: **green**

Flower symmetry: **actinomorphic**  
Perianth type: **homochlamydeous**  
Perianth fusion: **free**  
Inflorescence type: **spica**  
Dicliny: **synoecious**  
Generative reproduction type: **facultative autogamy**  
Pollination syndrome: **wind-pollination, water-pollination**

### **Fruit, seed and dispersal**

Fruit type: **dry fruit - achene/cypsela/samara**  
Fruit colour: **green, brown**  
Reproduction type: **by seed/spores and vegetatively**  
Dispersal unit (diaspore): **fruit, infrutescence or its part, turion, shoot fragment**  
Dispersal strategy: **Wolffia (mainly hydrochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot metamorphosis: **turion**  
Storage organ: **turion**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Karyology**

Chromosome number (2n): **26 (39)**  
Ploidy level (x): **2 (3)**  
2C genome size [Mbp]: **614.74**  
1Cx monoploid genome size [Mbp]: **307.37**  
Genomic GC content: **40.2 %**

### **Taxon origin**

Origin in the Czech Republic: **native**

### **Ecological indicator values**

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **12 - permanently or almost permanently submerged aquatic plant**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in**

### calcium-rich conditions

Nutrient indicator value: **6x - transition between values 5 and 7 (generalist)**

Salinity indicator value: **2 - oligohaline, often on soils with very low salt content**

### Habitat and sociology

Occurrence in habitats

3 Aquatic vegetation

3A Macrophytic vegetation of eutrophic and mesotrophic still waters: **2 - optimum**

3B Macrophytic vegetation of water streams: **1 - rare occurrence**

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 - rare occurrence**

4C Eutrophic vegetation of muddy substrata: **1 - rare occurrence**

4E Reed vegetation of brooks: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of associations: [VBB14 Parvo-Potamo-Zannichellietum pedicellatae](#), [VBB16 Najadetum marinae](#), [VBB17 Najadetum minoris](#)

Constant taxon

Constant taxon of associations: [VBB17 Najadetum minoris](#)

Dominant taxon

Dominant taxon of associations: [MCC09 Batrachio circinati-Alismatetum graminei](#), [VBB14 Parvo-Potamo-Zannichellietum pedicellatae](#), [VCA02 Charetum braunii](#)

### Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional, subtropical**

Floristic region: **Africa, Americas, circumpolar**

Distribution range extension along the continentality gradient: **9**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 384

taxon.data.freq\_in\_quad: 698

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**

Number of narrow habitats in which the taxon has its optimum: **1**

Number of broad habitats in which the taxon occurs: **2**

Number of broad habitats in which the taxon has its optimum: **1**

### Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**