


# *Potentilla thuringiaca*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



## Habitus and growth type

Height [m]: **0.15-0.5**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate, rosulate**

Leaf shape: **compound - ternate, compound - palmate (5-foliate)**

Stipules: **present**

Petiole: **present**

Leaf life span: **summer green**

## Flower

Flowering period [month]: **May-July**

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**

Flower colour: **yellow**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **aposepalous**  
Inflorescence type: **anthella**  
Dicliny: **synoecious**  
Generative reproduction type: **apomixis**  
Pollination syndrome: **insect-pollination**  
Pollinator spectrum: **solitary bees, flies s. l., other pollinators (honeybee, hoverflies, other Diptera, butterflies, beetles)**



## **Fruit, seed and dispersal**

Fruit type: **dry fruit - head of achenes**  
Reproduction type: **only by seed/spores**  
Dispersal unit (diaspore): **fruit, infrutescence or its part**  
Dispersal strategy: **Allium (mainly autochory)**  
Myrmecochory: **myrmecochorous nv**

## **Belowground organs and clonality**

Shoot metamorphosis: **pleiocorm**  
Storage organ: **pleiocorm**  
Type of clonal growth organ: **epigeogenous rhizome**  
Freely dispersible organs of clonal growth: **absent**  
Shoot life span (cyclicity): **monocyclic shoots prevailing**  
Branching type of stem-derived organs of clonal growth: **monopodial**  
Primary root: **absent**  
Persistence of the clonal growth organ [year]: **4**  
Number of clonal offspring: **0.5**  
Lateral spreading distance by clonal growth [m]: **0.01**  
Clonal index: **2**

### **Bud bank**

Number of buds per shoot at the soil surface (root buds excluded): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
Size of the belowground bud bank (root buds excluded): **20**  
Depth of the belowground bud bank (root buds excluded) [cm]: **4**  
Number of buds per shoot at the soil surface (root buds included): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
Size of the belowground bud bank (root buds included): **20**  
Depth of the belowground bud bank (root buds included) [cm]: **4**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in the Czech Republic: **native**

## Habitat and sociology

Occurrence in habitats

13 Anthropogenic vegetation

13F Herbaceous vegetation of forests clearings and Rubus scrub: **2 - optimum**

Affinity to the forest environment

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 - taxon occurring partly in the forest, but mainly in open vegetation**

Colonization ability

Index of colonization success (ICS): **1**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe, Western Asia**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 20

taxon.data.freq\_in\_quad: 27

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **1**

Number of narrow habitats in which the taxon has its optimum: **1**

Number of broad habitats in which the taxon occurs: **1**

Number of broad habitats in which the taxon has its optimum: **1**

## Threats and protection

Red List 2017 (national categories): **C1t - critically threatened taxon, declining**

Red List 2017 (IUCN categories): **CR - critically endangered**

Legal protection: **vulnerable taxon**