


Pteridium aquilinum

Distribution



© Dana Mjstřálcová

Map info

● revised records

● unrevised records


On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Petr Voborní



© Dana Mjstřálcová



© Eva Hřtenbergerová

Habitus and growth type

Height [m]: **0.5-2**

Growth form: **clonal herb**

Life form: **geophyte**

Life strategy: **C - competitor**

Life strategy (Pierce method based on leaf traits): **C**

Life strategy (Pierce method, C-score): **80.9 %**

Life strategy (Pierce method, S-score): **13.4 %**

Life strategy (Pierce method, R-score): **5.8 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **compound - bipinnate, compound - tripinnate**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **scleromorphic**

Flower

Flowering period [month]: **July-September**

Dicliny: **synoecious**

Fruit, seed and dispersal

Reproduction type: **by seed/spores and vegetatively**

Dispersal unit (diaspore): **spore**

Dispersal strategy: **Lycopodium (mainly anemochory)**

Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **rhizome**

Storage organ: **rhizome**

Type of clonal growth organ: **hypogeogenous rhizome**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicality): **monocyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **dichotomous**

Primary root: **absent**

Persistence of the clonal growth organ [year]: **4**

Number of clonal offspring: **1**

Lateral spreading distance by clonal growth [m]: **0.4**

Clonal index: **5**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **12**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **15**

Size of the belowground bud bank (root buds excluded): **27**

Depth of the belowground bud bank (root buds excluded) [cm]: **11**

Number of buds per shoot at the soil surface (root buds included): **0**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **12**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**

Size of the belowground bud bank (root buds included): **27**

Depth of the belowground bud bank (root buds included) [cm]: **11**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology


Chromosome number (2n): **104**

Ploidy level (x): **4**

2C genome size [Mbp]: **13397.43**

1Cx monoploid genome size [Mbp]: **3349.36**

Genomic GC content: **45.7 %**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **5 - indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **3 - acidity indicator, occurring mainly in acidic conditions, exceptionally in neutral conditions**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.83**

Herb layer disturbance frequency indicator value: **-0.62**

Whole-community disturbance severity indicator value: **0.38**

Herb layer disturbance severity indicator value: **0.26**

Whole-community structure based disturbance indicator value: **0.12**

Herb layer structure-based disturbance indicator value: **0.13**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**

6 Meadows and mesic pastures

6F Intermittently wet Molinia meadows: **1 - rare occurrence**

6G Vegetation of wet disturbed soils: **1 - rare occurrence**

7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 - rare occurrence**

8 Dry grasslands

8F Thermophilous forest fringe vegetation: **1 - rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 - rare occurrence**

9C Festuca grasslands on acidic sands: **1 - rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 - rare occurrence**

12 Forests

12A Alder carrs: **1 - rare occurrence**

12C Oak-hornbeam forests: **1 - rare occurrence**

12E Herb-rich beech forests: **1 - rare occurrence**

- 12G Acidophilous beech forests: **2 - optimum**
12I Sub-continental thermophilous oak forests: **1 - rare occurrence**
12K Acidophilous oak forests: **3 - dominant**
12L Boreo-continental pine forests: **2 - optimum**
12P Peatland pine forests: **1 - rare occurrence**
12Q Peatland birch forests: **2 - optimum**
12R Acidophilous spruce forests: **1 - rare occurrence**
12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**
12V Spruce plantations: **1 - rare occurrence**
12W Pine and larch plantations: **2 - optimum**

13 Anthropogenic vegetation

- 13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **3 - dominant**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.2 - taxon occurring mainly along forest edges and in forest openings, including forest roads and paths, windthrow sites, burnt sites and forest clearings**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.2 - taxon occurring mainly along forest edges and in forest openings, including forest roads and paths, windthrow sites, burnt sites and forest clearings**

Diagnostic taxon

Diagnostic taxon of alliances: [LFA Festuco-Pinion sylvestris](#), [LFB Dicrano-Pinion sylvestris](#)

Diagnostic taxon of associations: [LFA01 Festuco-Pinetum sylvestris](#), [XEA06 Pteridietum aquilini](#)

Constant taxon

Constant taxon of associations: [XEA06 Pteridietum aquilini](#)

Dominant taxon

Dominant taxon of associations: [XEA06 Pteridietum aquilini](#)

Ecological specialization indices

- Ecological specialization index for all vegetation types: **4.3**
Ecological specialization index for non-forest vegetation: **3.7**
Ecological specialization index for forest vegetation: **4.5**

Colonization ability

- Index of colonization success (ICS): **6**
Index of colonization potential (ICP): **9**
Optimum successional age [years]: **9**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional, subtropical, tropical, austral or antarctic**

Floristic region: **circumpolar**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**

Expansive taxon in the region: **Bohemian Moravian Mesophyticum, Bohemian Moravian Oreophyticum, Carpathian Mesophyticum, Carpathian Oreophyticum**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 536

taxon.data.freq_in_quad: 1367

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.3 %**

Occurrence frequency in vegetation plots with a cover above 5%: **50 %**

Occurrence frequency in vegetation plots with a cover above 25%: **33.3 %**

Occurrence frequency in vegetation plots with a cover above 50%: **23.3 %**

Mean percentage cover in vegetation plots: **25.5 %**

Maximum percentage cover in vegetation plots: **99 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **23**

Number of narrow habitats in which the taxon has its optimum: **6**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **2**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**