


# Rapistrum perenne

## Distribution



© Vladimír Nejšlechta

### Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Aleš Moravec

## Habitus and growth type

Height [m]: **0.3-0.8**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **C - competitor**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - pinnately divided**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **June-August**

Flowering phase: **7 Ligustrum vulgare-Stachys sylvatica (end of early summer)**

Flower colour: **yellow**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**  
Calyx fusion: **aposepalous**  
Inflorescence type: **racemus**  
Dicliny: **synoecious**  
Pollination syndrome: **insect-pollination**

### **Fruit, seed and dispersal**

Fruit type: **dry fruit - lomentum/transversely dehiscent siliqua**  
Reproduction type: **only by seed/spores**  
Dispersal unit (diaspore): **fruit, infrutescence or its part, tumbleweed**  
Dispersal strategy: **Allium (mainly autochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot metamorphosis: **pleiocorm**  
Root metamorphosis: **primary storage root**  
Storage organ: **pleiocorm, primary storage root**  
Shoot life span (cyclicity): **monocyclic shoots prevailing**  
Branching type of stem-derived organs of clonal growth: **sympodial**  
Primary root: **present**

#### **Bud bank**

Number of buds per shoot at the soil surface (root buds excluded): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
Size of the belowground bud bank (root buds excluded): **20**  
Depth of the belowground bud bank (root buds excluded) [cm]: **4**  
Number of buds per shoot at the soil surface (root buds included): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
Size of the belowground bud bank (root buds included): **20**  
Depth of the belowground bud bank (root buds included) [cm]: **4**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Karyology**

Chromosome number (2n): **18**  
Ploidy level (x): **2**  
2C genome size [Mbp]: **1333.3**  
1Cx monoploid genome size [Mbp]: **666.65**  
Genomic GC content: **40.2 %**

## Taxon origin

Origin in the Czech Republic: **native**

## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **7 - heat indicator, occurring in relatively warm lowlands**

Moisture indicator value: **3 - missing on damp soil**

Reaction indicator value: **8 - transition between values 7 and 9, occurring mostly in calcium-rich conditions**

Nutrient indicator value: **5 - occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.89**

Herb layer disturbance frequency indicator value: **-0.51**

Whole-community disturbance severity indicator value: **0.28**

Herb layer disturbance severity indicator value: **0.3**

Whole-community structure based disturbance indicator value: **0.76**

Herb layer structure-based disturbance indicator value: **0.7**

## Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8C Narrow-leaved sub-continental steppes: **2 - optimum**

8D Broad-leaved dry grasslands: **2 - optimum**

11 Heathlands and scrub

11N Low xeric scrub: **1 - rare occurrence**

13 Anthropogenic vegetation

13D Perennial thermophilous ruderal vegetation: **2 - optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Ecological specialization indices

Ecological specialization index for all vegetation types: **5**

Ecological specialization index for non-forest vegetation: **5**

Colonization ability

Index of colonization success (ICS): **4**

Index of colonization potential (ICP): **7**

Optimum successional age [years]: **75**

## Distribution and frequency

Floristic zone: **southern temperate, submeridional**

Floristic region: **Europe**

Continentality degree: **8**

Distribution range extension along the continentality gradient: **3**

Elevational belt in the Czech Republic: **lowlands, colline belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **58**

taxon.data.freq\_in\_quad: **113**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**

Occurrence frequency in vegetation plots with a cover above 5%: **0 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **1.9 %**

Maximum percentage cover in vegetation plots: **3 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **4**

Number of narrow habitats in which the taxon has its optimum: **3**

Number of broad habitats in which the taxon occurs: **3**

Number of broad habitats in which the taxon has its optimum: **2**

## Threats and protection

Red List 2017 (national categories): **C3 - vulnerable taxon**

Red List 2017 (IUCN categories): **NT - near threatened**

Legal protection: **not protected by law**