



Rumex obtusifolius

Distribution



© Dana Michalčová

Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Dana Michalčová



© Dana Michalčová



© Dana Michalčová

Habitus and growth type

Height [m]: **0.5-1.2**

Growth form: **polycarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **C - competitor**

Life strategy (Pierce method based on leaf traits): **C/CR**

Life strategy (Pierce method, C-score): **80.6 %**

Life strategy (Pierce method, S-score): **0 %**

Life strategy (Pierce method, R-score): **19.5 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **present**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic, hygromorphic**

Flower

Flowering period [month]: **June-August**

Flowering phase: **7 Ligustrum vulgare-Stachys sylvatica (end of early summer)**
 Flower colour: **green**
 Flower symmetry: **actinomorphic**
 Perianth type: **homochlamydeous**
 Perianth fusion: **free**
 Inflorescence type: **panicula e pseudospicis composita**
 Dicliny: **synoecious, gynomonoecious, androdioecious**
 Generative reproduction type: **facultative allogamy**
 Pollination syndrome: **wind-pollination**
 Pollinator spectrum: **hoverflies, other Diptera, beetles (other Hymenoptera, flies s. l., meat flies s. l.)**



Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**
 Fruit colour: **brown**
 Reproduction type: **mostly by seed/spores, rarely vegetatively**
 Dispersal unit (diaspore): **fruit, infructescence or its part**
 Dispersal strategy: **Allium (mainly autochory)**
 Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Shoot metamorphosis: **rhizome-like pleiocorm**
 Root metamorphosis: **primary storage root**
 Storage organ: **rhizome-like pleiocorm, primary storage root**
 Shoot life span (cyclicity): **monocyclic shoots prevailing**
 Branching type of stem-derived organs of clonal growth: **sympodial**
 Primary root: **present**
 Position of root buds: **primary root**
 Role of root buds in life-history of a plant: **regenerative**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **6**
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
 Size of the belowground bud bank (root buds excluded): **21**
 Depth of the belowground bud bank (root buds excluded) [cm]: **4**
 Number of buds per shoot at the soil surface (root buds included): **6**
 Number of buds per shoot at a depth of 0–10 cm (root buds included): **17**
 Number of buds per shoot at a depth greater than 10 cm (root buds included): **10**
 Size of the belowground bud bank (root buds included): **33**
 Depth of the belowground bud bank (root buds included) [cm]: **7**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
 Carnivory: **non-carnivorous**
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **40**

Ploidy level (x): **4**

2C genome size [Mbp]: **2272.71**

1Cx monoploid genome size [Mbp]: **568.18**

Genomic GC content: **40.9 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 - half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **6 - transition between values 5 and 7**

Reaction indicator value: **6x - transition between values 5 and 7 (generalist)**

Nutrient indicator value: **8 - pronounced nutrient indicator**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.83**

Herb layer disturbance frequency indicator value: **-0.38**

Whole-community disturbance severity indicator value: **0.57**

Herb layer disturbance severity indicator value: **0.57**

Whole-community structure based disturbance indicator value: **0.48**

Herb layer structure-based disturbance indicator value: **0.55**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 - rare occurrence**

4B Halophilous reed and sedge beds: **1 - rare occurrence**

4C Eutrophic vegetation of muddy substrata: **1 - rare occurrence**

4D Riverine reed vegetation: **2 - optimum**

4E Reed vegetation of brooks: **1 - rare occurrence**

4G Tall-sedge beds: **1 - rare occurrence**

4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**

4I Vegetation of nitrophilous annual hygrophilous herbs: **2 - optimum**

4J River gravel banks: **1 - rare occurrence**

4K Petasites fringes of montane brooks: **2 - optimum**

4L Nitrophilous herbaceous fringes of lowland rivers: **2 - optimum**

5 Vegetation of springs and mires

- 5A Hard-water springs with tufa formation: **1 - rare occurrence**
5B Lowland to montane soft-water springs: **1 - rare occurrence**
- 6 Meadows and mesic pastures
6A Mesic Arrhenatherum meadows: **1 - rare occurrence**
6B Montane mesic meadows: **1 - rare occurrence**
6C Pastures and park grasslands: **1 - rare occurrence**
6D Alluvial meadows of lowland rivers: **2 - optimum**
6E Wet Cirsium meadows: **1 - rare occurrence**
6F Intermittently wet Molinia meadows: **1 - rare occurrence**
6G Vegetation of wet disturbed soils: **2 - optimum**
- 7 Acidophilous grasslands
7A Subalpine and montane acidophilous grasslands: **1 - rare occurrence**
7B Submontane Nardus grasslands: **1 - rare occurrence**
- 10 Saline vegetation
10I Inland saline meadows: **1 - rare occurrence**
- 11 Heathlands and scrub
11I Willow carrs: **1 - rare occurrence**
11J Willow galleries of loamy and sandy river banks: **2 - optimum**
11L Tall mesic and xeric shrub: **1 - rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 - rare occurrence**
- 12 Forests
12A Alder carrs: **1 - rare occurrence**
12B Alluvial forests: **2 - optimum**
12T Robinia pseudacacia plantations: **1 - rare occurrence**
12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**
12V Spruce plantations: **1 - rare occurrence**
12W Pine and larch plantations: **1 - rare occurrence**
- 13 Anthropogenic vegetation
13A Annual vegetation of ruderal habitats: **1 - rare occurrence**
13B Annual vegetation of arable land: **1 - rare occurrence**
13C Annual vegetation of trampled habitats: **1 - rare occurrence**
13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 - optimum**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**
- Affinity to the forest environment
Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**
- Diagnostic taxon
Diagnostic taxon of classes: [KA *Salicetea purpureae*](#)
Diagnostic taxon of alliances: [KAC *Salicion albae*](#)
- Constant taxon
Constant taxon of associations: [KAC02 *Salicetum fragilis*](#)
- Dominant taxon
Dominant taxon of associations: [XCE01 *Urtico urentis-Chenopodietum boni-henrici*](#),

XDF01 *Rumicetum alpini*

Ecological specialization indices

Ecological specialization index for all vegetation types: **3.8**

Ecological specialization index for non-forest vegetation: **3.9**

Ecological specialization index for forest vegetation: **4.9**

Colonization ability

Index of colonization success (ICS): **6**

Index of colonization potential (ICP): **4**

Optimum successional age [years]: **7.5**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**

Expansive taxon in the region: **Bohemian Thermophyticum, Bohemian Moravian Mesophyticum, Pannonian Thermophyticum, Carpathian Mesophyticum**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **670**

taxon.data.freq_in_quad: **2403**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **4.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **7.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **1.4 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0.4 %**

Mean percentage cover in vegetation plots: **3.3 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **39**

Number of narrow habitats in which the taxon has its optimum: **9**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **5**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**