Sedum reflexum

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.1–0.25**
Growth form: **clonal herb**
Life form: **chamaephyte**
Life strategy: **S – stress-tolerator**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **evergreen**
Leaf anatomy: **succulent**

Flower

Flowering period [month]: **June–July**
Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **yellow**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Perianth fusion: **free**
Calyx fusion: **fused at the base**
Inflorescence type: **panicula e cincinnis composita**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – cluster of follicles**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed, shoot fragment**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably myrmecochorous nv**

Belowground organs and clonality

Shoot metamorphosis: **stolon, shoot succulence**
Storage organ: **stolon, succulence**
Type of clonal growth organ: **stolon**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **6**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **4**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **15**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **15**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **106 (88, 112)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ploidy level (x): **14 (12)**
2C genome size [Mbp]: **4986.56**
1Cx monoploid genome size [Mbp]: **356.18**
Genomic GC content: **42.1 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **2 – transition between values 1 and 3**
Reaction indicator value: **5 – indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions**
Nutrient indicator value: **2 – transition between values 1 and 3**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.3**
Herb layer disturbance frequency indicator value: **-0.62**
Whole-community disturbance severity indicator value: **0.23**
Herb layer disturbance severity indicator value: **0.23**
Whole-community structure based disturbance indicator value: **0.41**
Herb layer structure-based disturbance indicator value: **0.38**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**
1B Siliceous cliffs and block fields: **2 – optimum**
1D Mobile calcareous screes: **2 – optimum**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8B Submediterranean dry grasslands on rock outcrops: **2 – optimum**
8C Narrow-leaved sub-continental steppes: **2 – optimum**
8E Acidophilous dry grasslands: **2 – optimum**
8F Thermophilous forest fringe vegetation: **2 – optimum**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 – rare occurrence**
9C Festuca grasslands on acidic sands: **1 – rare occurrence**
9E Acidophilous vegetation of spring therophytes and succulents: **2 – optimum**
9F Basiphilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12J Acidophilous thermophilous oak forests: **2 – optimum**
12K Acidophilous oak forests: **1 – rare occurrence**
12L Boreo-continental pine forests: **1 – rare occurrence**
12T Robinia pseudacacia plantations: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of associations: **LCC02 *Genisto pilosae-Quercetum petraeae*, THA04 *Helichryso arenarii-Festucetum pallentis*, THC03 *Saxifrago paniculatae-Seslerietum caeruleae*, THG03 *Viscario vulgaris-Avenuletum pratensis***

Constant taxon

Constant taxon of associations: **LCC02 *Genisto pilosae-Quercetum petraeae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.5**
Ecological specialization index for non-forest vegetation: **4.5**
Ecological specialization index for forest vegetation: **5.3**

Colonization ability

Index of colonization success (ICS): **1**
Index of colonization potential (ICP): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **colline belt (lowlands)**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 217
taxon.data.freq_in_quad: 424

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.6 %**
Occurrence frequency in vegetation plots with a cover above 5%: **5 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0.6 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.7 %**
Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **20**
Number of narrow habitats in which the taxon has its optimum: **9**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz