

# *Sisymbrium austriacum*

## Distribution



© Dana Michalčová

### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Dana Michalčová



© Dana Michalčová

## Habitus and growth type

Height [m]: **0.3-0.6**

Growth form: **monocarpic perennial non-clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - pinnately divided**

Stipules: **absent**

Petiole: **mainly present**

Leaf life span: **evergreen**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **May-June**

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**

Flower colour: **yellow**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **free**  
 Calyx fusion: **aposepalous**  
 Inflorescence type: **panicula**  
 Dicliny: **synoecious**  
 Generative reproduction type: **mixed mating**  
 Pollination syndrome: **insect-pollination, selfing**

### Fruit, seed and dispersal

Fruit type: **dry fruit - siliqua**  
 Reproduction type: **only by seed/spores**  
 Dispersal unit (diaspore): **seed**  
 Dispersal strategy: **Allium (mainly autochory)**  
 Myrmecochory: **non-myrmecochorous (b)**

### Belowground organs and clonality

Root metamorphosis: **primary storage root**  
 Storage organ: **primary storage root**  
 Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**  
 Primary root: **present**

#### Bud bank

Number of buds per shoot at the soil surface (root buds excluded):  
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded):  
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded):  
 Size of the belowground bud bank (root buds excluded):  
 Depth of the belowground bud bank (root buds excluded) [cm]:  
 Number of buds per shoot at the soil surface (root buds included):  
 Number of buds per shoot at a depth of 0–10 cm (root buds included):  
 Number of buds per shoot at a depth greater than 10 cm (root buds included):  
 Size of the belowground bud bank (root buds included):  
 Depth of the belowground bud bank (root buds included) [cm]:

### Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**  
 Carnivory: **non-carnivorous**  
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in the Czech Republic: **neophyte**  
 Invasion status: **casual**  
 Geographic origin: **Europe, Mediterranean**  
 Year of the first record in the wild: **1858**  
 Period of introduction: **Late Modern Period (1800–1950)**



## Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 0

taxon.data.freq\_in\_quad: 0

## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**