Taraxacum ruptifolium

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **simple – pinnately divided**
Stipules: **absent**
Petiole: **present**

Flower

Flower colour: **yellow**
Flower symmetry: **zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **ligulate**
Calyx fusion: **pappus**
Inflorescence type: **anthodium solitarium**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Myrmecochory: **non-myrmecochorous (b)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **C4b – data deficient taxon**
Red List 2017 (IUCN categories): **DD – data deficient**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz