
Trachyspermum ammi

Distribution



Habitus and growth type

Growth form: **annual herb**
Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – bipinnate**
Stipules: **absent**
Petiole: **present**

Flower

Flower colour: **white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx absent, corolla present**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **umbella composita**

Fruit, seed and dispersal

Fruit type: **dry fruit – cremocarp**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **anecophyte**
Year of the first record in the wild: **1903**
Period of introduction: **Late Modern Period (1800–1950)**
Introduction pathway: **unintentional – industry**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz