


Vallisneria spiralis

Distribution



© Petr Kouřecký

Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.2-0.8**Growth form: **clonal herb**Life form: **hydrophyte**Life strategy (Pierce method based on leaf traits): **CR**Life strategy (Pierce method, C-score): **41.1 %**Life strategy (Pierce method, S-score): **0 %**Life strategy (Pierce method, R-score): **58.9 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**Leaf arrangement (phyllotaxis): **rosulate**Leaf shape: **simple - entire**Stipules: **absent**Petiole: **absent**Leaf life span: **evergreen**Leaf anatomy: **hydromorphic**

Flower

Flowering period [month]: **June-September**Flower colour: **green-white**

Perianth type: **reduced**
Perianth fusion: **reduced**
Calyx fusion: **aposepalous**
Inflorescence type: **flores solitarii**
Dicliny: **dioecious**
Generative reproduction type: **allogamy**
Pollination syndrome: **water-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**
Reproduction type: **only vegetatively**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon**
Storage organ: **stolon**
Type of clonal growth organ: **stolon**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **1.3**
Number of clonal offspring: **3.5**
Lateral spreading distance by clonal growth [m]: **0.09**
Clonal index: **5**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **8**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded):
Number of buds per shoot at a depth greater than 10 cm (root buds excluded):
Size of the belowground bud bank (root buds excluded): **13**
Depth of the belowground bud bank (root buds excluded) [cm]: **3**
Number of buds per shoot at the soil surface (root buds included): **8**
Number of buds per shoot at a depth of 0–10 cm (root buds included):
Number of buds per shoot at a depth greater than 10 cm (root buds included):
Size of the belowground bud bank (root buds included): **13**
Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**

Geographic origin: **Asia**

Period of introduction: **Present (after 2000)**

Introduction pathway: **intentional - ornamental**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **submeridional, meridional, subtropical**

Floristic region: **circumpolar**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 5

taxon.data.freq_in_quad: 5

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**