


# *Carex nigra* subsp. *nigra*

## Distribution



Map info	
●	revised records
○	unrevised records
On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.	



## Habitus and growth type

Height [m]: **0.1-0.8**

Growth form: **clonal herb**

Life form: **hemicryptophyte (geophyte)**

Life strategy: **S - stress-tolerator**

Life strategy (Pierce method based on leaf traits): **S/SR**

Life strategy (Pierce method, C-score): **7 %**

Life strategy (Pierce method, S-score): **68.8 %**

Life strategy (Pierce method, R-score): **24.3 %**



## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **helomorphic**



## Flower

Flowering phase: **4 Fagus sylvatica-Galeobdolon (start of mid-spring)**

Flower colour: **black**

Perianth type: **flower achlamydeous**

Inflorescence type: **spica e spiculis composita**

Dicliny: **monoecious, andromonoecious**

Generative reproduction type: **allogamy self-incompatibility**

Pollination syndrome: **wind-pollination**



© Jiří Procházka

## Fruit, seed and dispersal

Fruit type: **dry fruit - nut enclosed in an utricle**

Fruit colour: **brown**

Reproduction type: **by seed/spores and vegetatively**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (a)**



© Dana Michalcová

## Belowground organs and clonality

Shoot metamorphosis: **stolon, rhizome**

Storage organ: **stolon, rhizome**

Type of clonal growth organ: **epigeogenous rhizome**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **absent**

Persistence of the clonal growth organ [year]:

Number of clonal offspring:

Lateral spreading distance by clonal growth [m]: **0.09**

Clonal index: **4**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded):

Number of buds per shoot at a depth of 0-10 cm (root buds excluded):

Number of buds per shoot at a depth greater than 10 cm (root buds excluded):

Size of the belowground bud bank (root buds excluded):

Depth of the belowground bud bank (root buds excluded) [cm]:

Number of buds per shoot at the soil surface (root buds included):

Number of buds per shoot at a depth of 0-10 cm (root buds included):

Number of buds per shoot at a depth greater than 10 cm (root buds included):

Size of the belowground bud bank (root buds included):

Depth of the belowground bud bank (root buds included) [cm]:



© Dana Michalcová



© Dana Michalcová

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Karyology

Chromosome number (2n): **84**

Ploidy level (x): **2**

2C genome size [Mbp]: **786.69**

1Cx monoploid genome size [Mbp]: **393.34**

Genomic GC content: **36.2 %**



## Taxon origin

Origin in the Czech Republic: **native**



## Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **5 - moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**

Moisture indicator value: **8 - transition between values 7 and 9**

Reaction indicator value: **3 - acidity indicator, occurring mainly in acidic conditions, exceptionally in neutral conditions**

Nutrient indicator value: **2 - transition between values 1 and 3**

Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**



## Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.92**

Herb layer disturbance frequency indicator value: **-0.71**

Whole-community disturbance severity indicator value: **0.21**

Herb layer disturbance severity indicator value: **0.23**

Whole-community structure based disturbance indicator value: **0.53**

Herb layer structure-based disturbance indicator value: **0.66**



## Habitat and sociology

### Occurrence in habitats

2 Alpine and subalpine grasslands

2A Alpine grasslands on siliceous bedrock: **1 - rare occurrence**

2B Subalpine tall-forb and tall-grass vegetation: **1 - rare occurrence**

### 3 Aquatic vegetation

3C Macrophytic vegetation of oligotrophic lakes and pools: **1 - rare occurrence**

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 - rare occurrence**

4E Reed vegetation of brooks: **1 - rare occurrence**

4F Mesotrophic vegetation of muddy substrata: **1 - rare occurrence**

4G Tall-sedge beds: **2 - optimum**

4H Vegetation of low annual hygrophilous herbs: **1 - rare occurrence**

4K Petasites fringes of montane brooks: **1 - rare occurrence**

## 5 Vegetation of springs and mires

- 5B Lowland to montane soft-water springs: **1 - rare occurrence**
- 5C Alpine and subalpine soft-water springs: **1 - rare occurrence**
- 5D Calcareous fens: **2 - optimum**
- 5E Acidic moss-rich fens and peatland meadows: **3 - dominant**
- 5F Transitional mires: **3 - dominant**
- 5G Raised bogs: **2 - optimum**
- 5H Wet peat soils and bog hollows: **1 - rare occurrence**

## 6 Meadows and mesic pastures

- 6A Mesic Arrhenatherum meadows: **1 - rare occurrence**
- 6B Montane mesic meadows: **1 - rare occurrence**
- 6C Pastures and park grasslands: **1 - rare occurrence**
- 6D Alluvial meadows of lowland rivers: **2 - optimum**
- 6E Wet Cirsium meadows: **2 - optimum**
- 6F Intermittently wet Molinia meadows: **2 - optimum**
- 6G Vegetation of wet disturbed soils: **1 - rare occurrence**

## 7 Acidophilous grasslands

- 7A Subalpine and montane acidophilous grasslands: **1 - rare occurrence**
- 7B Submontane Nardus grasslands: **2 - optimum**

## 10 Saline vegetation

- 10I Inland saline meadows: **1 - rare occurrence**
- 10J Saline steppes: **1 - rare occurrence**
- 11 Heathlands and scrub
- 11A Dry lowland to subalpine heathlands: **1 - rare occurrence**
- 11D Subalpine acidophilous Pinus mugo scrub: **1 - rare occurrence**
- 11H Subalpine deciduous scrub: **1 - rare occurrence**
- 11I Willow carrs: **1 - rare occurrence**
- 11J Willow galleries of loamy and sandy river banks: **1 - rare occurrence**

## 12 Forests

- 12A Alder carrs: **1 - rare occurrence**
- 12G Acidophilous beech forests: **1 - rare occurrence**
- 12K Acidophilous oak forests: **1 - rare occurrence**
- 12P Peatland pine forests: **1 - rare occurrence**
- 12Q Peatland birch forests: **2 - optimum**
- 12R Acidophilous spruce forests: **1 - rare occurrence**
- 12S Basiphilous spruce forests: **1 - rare occurrence**
- 12V Spruce plantations: **1 - rare occurrence**

## 13 Anthropogenic vegetation

- 13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 - rare occurrence**

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 - taxon occurring partly in the forest, but mainly in open vegetation**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 - taxon occurring partly in the forest, but mainly in open vegetation**

### Diagnostic taxon

Diagnostic taxon of classes: [\*\*RB Scheuchzerio palustris-Caricetea nigrae\*\*](#)

Diagnostic taxon of alliances: [RBA Caricion davallianae](#), [RBB Sphagno warnstorpii-Tomentypnion nitentis](#), [RBC Caricion canescens-nigrae](#), [RBD Sphagno-Caricion canescens](#)

Diagnostic taxon of associations: [RBA03 Valeriano simplicifoliae-Caricetum flavae](#), [RBB01 Sphagno warnstorpii-Eriophoretum latifolii](#), [RBC01 Caricetum nigrae](#), [RBC03 Agrostio caninae-Caricetum diandrae](#), [RBD03 Carici echinatae-Sphagnetum](#), [TDF03 Angelico sylvestris-Cirsietum palustris](#)

Constant taxon

Constant taxon of classes: [RB Scheuchzerio palustris-Caricetea nigrae](#)

Constant taxon of alliances: [RBA Caricion davallianae](#), [RBC Caricion canescens-nigrae](#), [RBD Sphagno-Caricion canescens](#), [TDD Molinion caeruleae](#), [TDF Calthion palustris](#), [TED Nardo strictae-Juncion squarroso](#)

Constant taxon of associations: [LAA01 Thelypterido palustris-Alnetum glutinosae](#), [LAB01 Salicetum auritae](#), [MCG04 Comaro palustris-Caricetum cespitosae](#), [MCG06 Caricetum appropinquatae](#), [RBA01 Valeriano dioicae-Caricetum davallianae](#), [RBA02 Carici flavae-Cratoneuretum filicini](#), [RBA03 Valeriano simplicifoliae-Caricetum flavae](#), [RBA04 Campylio stellati-Caricetum lasiocarpae](#), [RBB01 Sphagno warnstorpii-Eriophoretum latifolii](#), [RBB02 Campylio stellati-Trichophoretum alpini](#), [RBB03 Menyantho trifoliatae-Sphagnetum teretis](#), [RBC01 Caricetum nigrae](#), [RBC03 Agrostio caninae-Caricetum diandrae](#), [RBD01 Sphagno recurvi-Caricetum rostratae](#), [RBD02 Sphagno recurvi-Caricetum lasiocarpae](#), [RBD03 Carici echinatae-Sphagnetum](#), [TDD02 Junco effusi-Molinietum caeruleae](#), [TDF01 Angelico sylvestris-Cirsietum oleracei](#), [TDF02 Cirsietum rivularis](#), [TDF03 Angelico sylvestris-Cirsietum palustris](#), [TDF04 Crepido paludosae-Juncetum acutiflori](#), [TDF06 Chaerophyllo hirsuti-Calthetum palustris](#), [TDF08 Scirpetum sylvatici](#), [TDF10 Scirpo sylvatici-Caricetum brizoidis](#), [TED01 Juncetum squarroso](#)

Dominant taxon

Dominant taxon of associations: [RBA03 Valeriano simplicifoliae-Caricetum flavae](#), [RBC01 Caricetum nigrae](#), [RBD03 Carici echinatae-Sphagnetum](#), [TDF03 Angelico sylvestris-Cirsietum palustris](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.5**

Ecological specialization index for non-forest vegetation: **5.6**

Ecological specialization index for forest vegetation: **5.1**

Colonization ability

Index of colonization success (ICS): **6**

Index of colonization potential (ICP): **2**

Optimum successional age [years]: **28**

## Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Asia, Americas**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 622

taxon.data.freq\_in\_quad: 1942

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **5.4 %**

Occurrence frequency in vegetation plots with a cover above 5%: **26.5 %**

Occurrence frequency in vegetation plots with a cover above 25%: **8.2 %**

Occurrence frequency in vegetation plots with a cover above 50%: **2.1 %**

Mean percentage cover in vegetation plots: **7.8 %**

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **41**

Number of narrow habitats in which the taxon has its optimum: **10**

Number of broad habitats in which the taxon occurs: **10**

Number of broad habitats in which the taxon has its optimum: **5**

## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**