


# *Datura stramonium* var. *tatula*

## Distribution



© Veronika Kahlíková

### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Veronika Kahlíková



© Veronika Kahlíková

## Habitus and growth type

Life form: **therophyte**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**

## Flower

Flower colour: **violet**

Flower symmetry: **actinomorphic**

Perianth type: **calyx and corolla**

Perianth fusion: **fused**

Shape of the sympetalous corolla or syntepalous perianth: **funnel-shaped**

Calyx fusion: **synsepalous**

Inflorescence type: **flores solitarii**

## Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**

Dispersal unit (diaspore): **seed**  
Dispersal strategy: **Allium (mainly autochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Taxon origin**

Origin in the Czech Republic: **neophyte**  
Invasion status: **naturalized**  
Geographic origin: **North America**  
Year of the first record in the wild: **1934**  
Period of introduction: **Late Modern Period (1800-1950)**  
Introduction pathway: **intentional - ornamental, unintentional - agriculture, unintentional - nature**

### **Ecological indicator values**

#### Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**  
Temperature indicator value: **7 - heat indicator, occurring in relatively warm lowlands**  
Moisture indicator value: **4 - transition between values 3 and 5**  
Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**  
Nutrient indicator value: **8 - pronounced nutrient indicator**  
Salinity indicator value: **1 - salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

### **Habitat and sociology**

#### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**  
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**  
Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**  
Legal protection: **not protected by law**